



Roloff, E. V. L., Tomiak-Baquero, A. M., Kasparov, S., & Paton, J. F. R. (2016). Parasympathetic innervation of vertebrobasilar arteries: is this a potential clinical target? *Journal of Physiology*, 594(22), 6463-6485. <https://doi.org/10.1113/JP272450>

Peer reviewed version

License (if available):
Unspecified

Link to published version (if available):
[10.1113/JP272450](https://doi.org/10.1113/JP272450)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via Wiley at <https://physoc.onlinelibrary.wiley.com/doi/abs/10.1113/JP272450> . Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Parasympathetic innervation of vertebro-basilar arteries: is this a potential clinical target?

Eva v. L. Roloff, Ana M. Tomiak-Baquero, Sergey Kasparov & Julian F.R. Paton

School of Physiology, Pharmacology & Neuroscience, Biomedical Sciences, University of
Bristol, Bristol BS8 1TD, UK.

Corresponding Author

Professor Julian F.R. Paton

School of Physiology, Pharmacology & Neuroscience,

Biomedical Sciences

University of Bristol

Bristol BS8 1TD, UK.

Julian.f.r.paton@Bristol.ac.uk

Running title: Vertebro-basilar artery innervation

Size: approximately 10 pages including figures, ~6988 words.

Summary

This review aims to summarise the contemporary evidence for the presence and function of the parasympathetic innervation of the cerebral circulation with emphasis on the vertebral and basilar arteries (the posterior cerebral circulation). We consider whether the parasympathetic innervation of blood vessels could be used as a means to increase cerebral blood flow. This may have clinical implications for pathologies associated with cerebral hypoperfusion such as for stroke, dementia and hypertension. Relative to the anterior cerebral circulation little is known of the origins and neurochemical phenotypes of the parasympathetic innervation of the vertebrobasilar arteries. These vessels normally provide blood flow to the brainstem and cerebellum, but can via the Circle of Willis upon stenosis of the internal carotid arteries supply blood to the anterior cerebral circulation too. We review the multiple types of parasympathetic fibres and their distinct transmitter mechanisms and how these vary with age, disease and species. We highlight the importance of parasympathetic fibres for mediating the vasodilatory response to sympathetic activation. Current trials are investigating the possibility of electrically stimulating the postganglionic parasympathetic ganglia to improve cerebral blood flow to reduce the penumbra following stroke. We conclude that although there are substantial gaps in our understanding of the origins of parasympathetic innervation of the vertebrobasilar arteries, activation of this system under some conditions might bring therapeutic benefits.

List of abbreviations

7-NI	7-nitroindazole
ACA	anterior cerebral artery
AChE	acetyl choline esterase
AcoA	anterior communicating artery
ADMA	asymmetric dimethylarginin
BBB	blood brain barrier
CBF	cerebral blood flow
CGRP	calcitonin gene related peptide
ChAT	choline-acetyl-transferase
CMG	carotid mini-ganglia
DBH	dopamine β -hydroxylase
ECA	external carotid artery
eNOS	endothelial nitric oxide synthase
ICA	internal carotid artery
iNOS	induceable nitric oxide synthase
L-NA	N ^G -nitro-L-arginine
NACHR	nicotinic acetylcholine receptor
NAPDH	nicotinamide adenine dinucleotide phosphate
nNOS	neuronal nitric oxide synthase
NO	nitric oxide
NOS	nitric oxide synthase
NPR-C	natriuretic peptide clearance receptor
NPY	neuropeptide Y
PACAP	pituitary adenylate cyclase-activating peptide
PCA	posterior cerebral artery
PCoA	posterior communicating artery
PGF _{2α}	prostaglandin F _{2α}
PHI	peptide histidine isoleucine
PPG	pterygopalatine ganglia
RT-PCR	real time polymerase chain reaction
SHR	spontaneously hypertensive rat
SMC	smooth muscle cell
SP	substance P
SPG	sphenopalatine ganglia
SSN	superior salivatory nucleus
TH	tyrosine hydroxylase
VACHT	vesicular acetyl choline transferase
VIP	vasoactive intestinal peptide
VPAC	VIP receptor
WGA-HRP	wheat germ agglutinin-horse radish peroxidase

Introduction

Innervation of the cerebral vasculature was described by Willis as early as 1664, but not until advances in histochemical and immunofluorescent techniques and electrophysiological and pharmacological investigations in the 20th century, did we begin to learn about its origins and function. Innervation of the anterior cerebral circulation, which supply the cerebral hemispheres and deliver the majority of blood to the brain, has been described in detail in several reviews (Purves, 1972); Dahl (1973); (Bevan *et al.*, 1981; Edvinsson, 1987; Bley & Cowen, 2001; Gulbenkian *et al.*, 2001; Lee *et al.*, 2001; Hamel, 2006; Owman, 2011; Goadsby, 2013). Our aim is to provide an up to date account of the parasympathetic innervation of the posterior vessels (e.g vertebral and basilar), which have received far less attention and whose normal physiological function is still largely unknown.

Clinically, there is a strong interest in cerebral blood flow regulation because of the devastating personal and socio-economic consequences of long term hypo-perfusion and stroke. As the brainstem controls the most essential brain functions for life, the consequences of reduced blood flow in this part of the brain are devastating. Thus, understanding the control of blood flow to the posterior circulation of the brain is critically important and any vasodilating mechanisms highly relevant clinically as it may provide a novel therapeutic target. In the context of ischemic stroke, the parasympathetic innervation to the vertebral and basilar arteries that supply blood flow to the brainstem (and cerebellum) are of great interest and have not been reviewed systematically.

Gross anatomy of cerebral arteries

The brain is one of the most metabolic demanding tissues in the body and it is not surprising therefore that the main cerebral arteries possess several specialisations that make them different from other arteries in the body enabling them to meet the high demand on blood flow. Their organisation is unique such that multiple feed arteries are interconnected with anastomoses forming a ring like structure: the Circle of Willis (Fig 1). The unique blood vessel organisation of the Circle of Willis acts as a distribution centre permitting blood to flow in any direction to meet increased demand and overcome stenosis. However, under normal circumstances blood flow distribution is normally well demarcated with little mixing from the different feeder arteries (McDonald & Potter, 1951). Thus, in the human and rat cerebral blood flow is mainly provided by the internal carotid arteries (ICA) whereas the vertebral and spinal arteries supply the brainstem.

-Insert Fig 1 here-

(human angiogram and schematic of cerebral circulation)

The Circle of Willis is formed from several inter-connecting arteries: including the posterior cerebral artery (PCA), posterior communicating artery (PCoA), anterior cerebral artery (ACA) and anterior communicating artery (AcoA); the latter does not exist in rats as the ACAs fuse to form the azygos anterior cerebral artery (Scremin, 2004). There is considerable variation to this organisation both inter- (Daniel *et al.*, 1953; McFarland *et al.*, 1979) and intra-species (Alpers *et al.*, 1959; Brown, 1966; Frąckowiak & Jakubowski, 2008), as seen in Figure 2. In humans, monkeys and rodents the ICAs supply blood to the Circle of Willis with minimal contribution from the external carotid artery (ECA). In most other mammals including cats, dogs, pigs and sheep, a large proportion of brain blood flow originates from the external

carotid artery via a rete mirabili¹ before joining the Circle of Willis with either no or a small contribution from the internal carotids (Daniel *et al.*, 1953). In humans, primates, rabbits and rats there is no rete.

-Insert Fig 2 here-

(Species variation in cerebral artery organisation)

A unique feature of cerebral arteries is that they are both conduit and resistance vessels and unlike other arteries constrict and dilate (Faraci & Heistad, 1990). Constriction can increase arterial pressure by ~40% in human (Dickinson & Paton, 2012) and rat (Baumbach & Heistad, 1983). This ability protects the brain from sudden rises in BP to prevent stroke (Bill & Linder, 1976; Edvinsson *et al.*, 1976a; Beausang-Linder & Bill, 1981; Sadoshima *et al.*, 1981), whereas vasodilation can protect against ischemia (Santizo *et al.*, 2000; Levi *et al.*, 2012). During constriction maintenance of blood flow to the brain then becomes dependent on compensatory vasodilatation in parenchymal arterioles (Baumbach & Heistad, 1983).

-Insert Figure 3 here-

(Arterial contribution to cerebral blood supply in different species)

¹ A rete mirabili is a complex of arteries and veins in close proximity and built on a counter-current principle, which creates a gradient that enables exchange of heat, ions or gas between the vessels. The function of the rete mirabili varies between species and is controversial in others. Barnett CH & Marsden CD. (1961). Functions of the mammalian carotid rete mirabile. *Nature* **191**, 88-89. It has been suggested it provide means to regulate the temperature of blood flowing to the brain in dogs, cats, sheep and goats. Baker MA. (1982). Brain cooling in endotherms in heat and exercise. *Annual review of physiology* **44**, 85-85. In the dolphin an extensive rete provide means of pressure dampening of BP in the brain. Nagel EL, Morgane PJ, McFarland WL & Galllano RE. (1968). Rete mirabile of dolphin: its pressure-damping effect on cerebral circulation. *Science* **161**, 898-900. A similar function has been attributed to the rete mirabile in the giraffe. Lawrence WE & Rewell RE. (1948). The cerebral blood supply in the giraffidae. *Proceedings of the Zoological Society of London* **118**, 202-212..

Neural innervation of major cerebral vessels

There are two sources of innervation controlling blood flow to cerebral vessels: extrinsic and intrinsic (Hamel, 2006). 'Intrinsic' innervation originating from local neurons within the CNS targeting arterioles to control blood flow in the brain parenchyma. Our focus is on the 'extrinsic' innervation from peripheral ganglia that innervate all major cerebral arteries up to the Virchow-Robin space², that is, before they enter the brain parenchyma.

Extrinsic innervation consists of the sensory system and efferent motor-fibres from the autonomic (sympathetic and parasympathetic) nervous system. Numerous proteins have been associated with functionally different subsets of nerve fibres that can be detected using immunocytochemistry. Examples relevant to the cerebral circulation include: acetylcholinesterase (AChE), vesicular acetylcholine transferase (VACHT) and vasoactive intestinal peptide (VIP) within parasympathetic fibres; tyrosine hydroxylase (TH), dopamine β -hydroxylase (DBH), and neuropeptide Y (NPY) in sympathetic nerves, and calcitonin gene related peptide (CGRP) as a marker of sensory afferents. In basilar arteries from human cadaveric fetuses both noradrenergic and AChE containing fibres first appear around the 12th gestational week whereas NPY and VIP synchronously appeared a month later at 16th gestational week (Kawamura & Takebayashi, 1994). Interestingly, with aging, there was a *decrease* in the expression of vasoconstrictor neurotransmitters (e.g. TH) in cerebrovascular nerves, whereas the expression of vasodilator neurotransmitters (VIP and CGRP) in

² Virchow Robin spaces are perivascular spaces or canals surrounding arteries and veins at the point where they leave the subarachnoid on the surface of brain and turn into the parenchyma. See figure in Hamel E. (2006). Perivascular nerves and the regulation of cerebrovascular tone. *Journal of applied physiology* **100**, 1059-1064.

perivascular nerve fibres supplying the rat cerebral arteries *increased* with age (Mione *et al.*, 1988).

Stimulation of sympathetic nerves is generally considered to result in vasoconstriction (D'Alecy & Feigl, 1972; Lee *et al.*, 1976; Baumbach & Heistad, 1983), whereas stimulation of parasympathetic nerves is vasodilatory (D'Alecy & Rose, 1977; Talman *et al.*, 2007). As we shall see later, there are exceptions to this rule. Sensory neurons are a heterogeneous population of either fast transmitting A δ fibres (being purely afferent) or slowly conducting capsaicin sensitive fibres expressing a range of co-transmitters. Exceptionally, these fibres can also affect cerebrovascular tone due to their additional efferent role; these fibres are referred to as “sensory-motor nerves” (Rubino & Burnstock, 1996) and contain, for example, substance P (SP) and calcitonin gene related peptide (CGRP), a very potent vasodilator.

Extrinsic nerve plexuses and bundles are located on the periphery of the adventitia, mostly run in the longitudinal direction along the vessel and have a smooth (‘string like’) appearance, whereas single fibres and terminals tend to be located in the deeper layers of the adventitia, running circumferential/perpendicular to the vessel and close to the smooth muscle cells (SMC) (Bleys *et al.*, 1996b; Taktakishvili *et al.*, 2010). The fine terminal nerve fibres are characterised by varicosities and have a ‘pearl-on-a string’ appearance (Yoshida *et al.*, 1993). These structures contain the highest vesicle numbers and neurotransmitter contents. Autonomic and sensory nerve fibres can be in close proximity to smooth muscle cells (80-100nm) and to each other (Iwayama *et al.*, 1970; Matsuyama *et al.*, 1985).

However, on basilar arteries close contacts between varicosities and smooth muscle cells are sparse and distances of 1-2 μ m are typical (Luff & McLachlan, 1989; Lee *et al.*, 1990). The

functional significance of this is unclear but it may reduce efficacy of transmission in these structures.

The parasympathetic innervation of posterior cerebral vessels

Cerebral arteries are richly innervated with neuronal fibres of parasympathetic origin providing a most powerful vasodilatory mechanism (Goadsby, 2013). Parasympathetic fibres follow the sympathetic nerves along cerebral vessels. In comparison to peripheral arteries (femoral and common carotid), where parasympathetic innervation is negligible, it is generally 10-40 times higher in cerebral arteries (Duckles, 1981). The density of innervation is both species and cerebral artery location dependent. Using vasoactive intestinal peptide (VIP) immunopositive fibres as a marker for parasympathetic innervation, relatively low densities have been observed in mouse, rabbit and monkey, with intermediate levels in dog and guinea pig, whereas pig, rat and bat exhibited high levels (Edvinsson *et al.*, 1980; Ando, 1988). In general, fibre densities are greater in the rostral part of the cerebral circulation, than in the caudal (Edvinsson *et al.*, 1980; Kobayashi *et al.*, 1983; Hara & Weir, 1986; Hara & Kobayashi, 1987). The exception is the bat, demonstrating dense VIP positive innervation of the vertebrobasilar arteries (Ando, 1988). In some animals (dogs, rabbits), either no or low parasympathetic innervation of the vertebrobasilar arteries has been reported (Florence & Bevan, 1979; Edvinsson *et al.*, 1980; Saito *et al.*, 1985). However, dogs have NOS positive fibre staining on vertebrobasilar arteries (Yoshida *et al.*, 1993) that was shown not to be sensory, suggesting the presence of some 'parasympathetic' fibres.

Types of parasympathetic fibres innervating posterior cerebral vessels

Classically, peripheral parasympathetic nerves are considered to be cholinergic, i.e. using acetylcholine (ACh) as a transmitter. However, it is clear that parasympathetic cells innervating cerebral arteries utilise a host of co-transmitters in addition or in preference to ACh. Some parasympathetic neurones in the cerebral arteries have lost the ability to produce ACh – thus giving rise to the term ‘non-cholinergic parasympathetic innervation’ (Minami *et al.*, 1994).

Nitric oxide (NO), vasoactive intestinal peptide (VIP), and/or pituitary adenylate cyclase-activating peptide (PACAP) have all been identified as co-transmitters in parasympathetic neurones and all are potent vasodilators (Bevan *et al.*, 1984; McCulloch *et al.*, 1986; Warren *et al.*, 1991; Nozaki *et al.*, 1993; Uddman *et al.*, 1999) producing greater vasodilatation than ACh itself (Lee, 2000). Reviewing the evidence of vasodilation in large cerebral arteries, Lee (2002) concluded that the main effector for vasodilation is NO, with the other neurotransmitters, including ACh, acting as presynaptic modulators of NO release. These findings explained the initial confusion that existed after cholinergic nerves had been identified on cerebral arteries, but were found to be resistant to atropine/physostigmine vasodilation, leading to the misconception that these nerves were non-functional (Lee *et al.*, 1978; Lee, 1980, 1982). Thus, parasympathetic neurons innervating the posterior cerebral arteries consist of a heterogeneous population of ACh-VIP-NO-, ACh-NO- and VIP-NO-positive neurons - but this list is not exhaustive (Minami *et al.*, 1994; Kimura *et al.*, 1997; Yu *et al.*, 1998). It is important to note, that NO and PACAP are also co-expressed in sensory neurones on cerebral arteries (Sundler *et al.*, 1996). Therefore, immunocytochemical labelling of parasympathetic neurones is challenging and requires co-labelling.

Tract tracing studies of parasympathetic input to the cerebral arteries have employed a variety of antibodies including cholinergic markers: choline-acetyl-transferase (ChAT) or the vesicular-acetylcholine-transporter (VACHT), or VIP and the NO producing enzymes: NO synthetase (NOS) or NADPH-diaphorase. Earlier immunohistochemical studies also used acetyl choline esterase (AChE)³, though this enzyme is less specific. AChE degrades ACh and was one of the first available cholinergic markers; however, it is not exclusive to cholinergic neurones and has been considered as a pan neuronal marker (Bleys *et al.*, 1996b).

The relative distribution of the different markers (ChAT, VACHT, NOS and VIP) and proportion of neurones expressing them has not been fully elucidated for the vertebrobasilar arteries and reports vary greatly partly due to species differences. ChAT has proven capricious in the periphery and in fine fibres due to low enzyme levels and hence its use underestimates the total cholinergic innervation (Schäfer *et al.*, 1998; Bleys *et al.*, 2001).

Origin of parasympathetic fibres to the posterior cerebral arteries

The origin of the parasympathetic innervation is characterised by a diffuse collection of small extra-cranial ganglia and small cell groups in both the arachnoid and at the base of the skull.

-Insert Figure 4 here-

(Overview of Parasympathetic innervation)

³ Acetylcholine esterase is an enzyme involved in the breakdown of ACh in the synaptic cleft. This was one of the first available cholinergic associated markers. In addition to cholinergic cells it also stains many adrenergic and sensory nerves (Bleys RL, Groen GJ & Hommersom RF. (1996a). Neural connections in and around the cavernous sinus in rat, with special reference to cerebrovascular innervation. *The Journal of comparative neurology* **369**, 277-291..

The anterior circulation is innervated by fibres from the pterygopalatine ganglia (PPG), also referred to as the sphenopalatine ganglia (SPG), the cavernous sinus, the otic and the internal carotid mini-ganglia (CMG) (Chorobski & Penfield, 1932; Kobayashi *et al.*, 1983; Carvalho, 1985; Keller *et al.*, 1985; Hara & Weir, 1986; Walters *et al.*, 1986; Hara & Kobayashi, 1987; Suzuki *et al.*, 1988; Edvinsson *et al.*, 1989; Hara *et al.*, 1989; Hardebo *et al.*, 1991; Suzuki & Hardebo, 1991; Nakai *et al.*, 1993; Suzuki & Hardebo, 1993; Bleys *et al.*, 1996a; Toda *et al.*, 2000a; Toda *et al.*, 2000b; Bleys *et al.*, 2001). The origin of parasympathetic innervation to the vertebrobasilar arteries is less well understood. In monkeys, PPG fibres appeared to be the source of NOS positive fibres to the anterior but not posterior cerebral arteries (Ayajiki *et al.*, 2012) and in rats parasympathetic VIP- (Hara *et al.*, 1989) and NOS- positive fibres (Nozaki *et al.*, 1993) were found on caudal basilar and vertebral arteries following PPG extirpation, suggesting additional sources. Injecting the anterograde tracer wheat germ agglutinin-horse radish peroxidase (WGA-HRP) into the PPG, only stained parasympathetic fibres at the anterior end of the basilar artery (Hara *et al.*, 1993). Similar injections to the otic ganglia did not result in labelling on the vertebrobasilar arteries in rats (Shimizu, 1994). However, using retrograde tract tracing with fluoro-gold applied to the middle portion of the basilar artery resulted in neurone labelling in both the otic ganglia and PPG (Kadota *et al.*, 1996). Labelling in the PPG was also seen following WGA-HRP application to the middle and caudal part of the BAs in cats (Keller *et al.*, 1985). Bleys and Cowen (2001) suggested that PPG, otic and CMG innervation is restricted to the anterior end of the basilar artery with no observable innervation to the caudal basilar and vertebral arteries. Gibbins *et al.* (1984a) lists otic, PPG, submandibular and sublingual ganglia as possible contributions to rostral basilar artery VIP containing fibres. The inconsistencies may be explained by the differences in techniques and species used. Studies

where NOS immunocytochemistry has been performed will not be able to delineate between parasympathetic and sensory nerves. Hence, many of the suggested sources of parasympathetic innervation to the posterior cerebral arteries based on NOS immunochemistry, which included the glossopharyngeal and vagal nerves, and upper cervical root ganglia (Suzuki *et al.*, 1988; Hardebo *et al.*, 1991), are all likely to be of sensory origin (Arbab *et al.*, 1986).

Despite evidence for parasympathetic innervation of vertebrobasilar arteries, and the existence of: VAcHT (E. Roloff & JFR Paton, personal observation), ChAT and VIP positive fibres in rat (Suzuki *et al.*, 1988; Suzuki *et al.*, 1990c), cat (Gibbins *et al.*, 1984b) and dog (Seki *et al.*, 1995), we have found sparse information regarding the source(s) of these specific parasympathetic inputs. This clearly needs further clarification.

Pre-ganglionic parasympathetic neurones destined for the cerebral vasculature

The location of preganglionic neurones projecting to the PPG of rats has been determined by retrograde transneuronal pseudorabies virus labelling (Spencer *et al.*, 1990). Neuronal labelling was obtained in the ipsilateral superior salivatory nucleus (SSN). Pharmacological stimulation of the greater petrosal nerve cell group within the SSN reduces ipsilateral cerebral vascular resistance. The effect can be countered by PPG section but not with systemic muscarinic antagonism; another confirmation that the vasodilatory mechanism (via the PPG) is mediated by a non-cholinergic mechanism in the cerebral arteries (Nakai *et al.*, 1993). The following nuclei were also labelled and most likely upstream of the SSN: the nucleus tractus solitarius and the dorso medial part of the spinal trigeminal nucleus and gigantocellular reticular nucleus as well as pontine structures of the parabrachial nucleus,

A5 catecholaminergic cells, and sub coeruleus region of the pons. Midbrain areas included the lateral, dorsomedial and paraventricular hypothalamic nuclei, lateral preoptic area. The bed nucleus of the stria terminalis, substantia innominata and the amygdalopiriform were also labelled in the forebrain. Given this extensive convergent connectivity it underpins the importance of the vasodilatory role of the SSN and those regions that may influence it.

There is scant information regarding sources of input to the other parasympathetic ganglia innervating cerebral arteries. Gibbins et al. (1984) and Gibbins & Morris (1988) reported the SSN was also a source of input to the otic ganglia and CMG.

Functional role of the parasympathetic transmitters and receptors

The transmitters and receptors are presented in order of importance as vasodilatory mechanisms.

(i) Nitrenergic: NO is the main vasodilator in the cerebral arteries (Lee, 2002). It has an estimated diffusion range of up to 300µm and an estimated half-life of 5-15s in the cerebral circulation (Ignarro, 1991; Garthwaite & Boulton, 1995). Nerve fibres with NO production capabilities on cerebral arteries include pure nitrenergic, VIPergic and/or cholinergic parasympathetic neurones as well as sensory neurones (Minami *et al.*, 1994; Kimura *et al.*, 1997; Edvinsson *et al.*, 2001). Note that there are multiple additional sources of NO that could produce vasodilation in the vertebrobasilar arteries such as endothelial cells, astrocytes, neurones (Bredt *et al.*, 1990; Faraci & Brian, 1994) and immune cells (Hibbs Jr *et al.*, 1988; Bogdan, 2001). The distribution of NO containing fibres in the cerebral arteries is highly variable in terms of exact location on the cerebral arteries and species. In rats, although present on the basilar artery, the densest NOS/NADPH-diaphorase positive fibre

staining occurred on the anterior part of the cerebral circulation, whereas in humans NOS immuno-reactivity was predominantly on posterior cerebral arteries, especially the basilar artery (Iadecola *et al.*, 1993; Nozaki *et al.*, 1993; Minami *et al.*, 1994; Edvinsson *et al.*, 2001; Taktakishvili *et al.*, 2010).

(ii) Cholinergic: Cholinergic receptors have been located on endothelial cells, smooth muscle cells, and perivascular nerves on cerebral vessels (see Table 1). Electrical or pharmacological stimulation of cholinergic fibres on cerebral arteries can, depending on the species, the site of action, receptor subtype and concentration of ACh released, induce either vasodilatation or vasoconstriction (Edvinsson *et al.*, 1977; Duckles, 1981; Miao & Lee, 1991; Dauphin & Mackenzie, 1995; Toda *et al.*, 1997). In cats and dogs, low concentrations of ACh (10^{-6} to 10^{-5} M) applied to cerebral arteries including the basilar caused vasodilation, whereas higher concentrations result in vasoconstriction (Tsukahara *et al.*, 1986; Dauphin & Hamel, 1992). In humans, however, only vasodilation was observed (Edvinsson *et al.*, 1976b; Tsukahara *et al.*, 1986). ACh can exert its effects via either metabotropic muscarinic (M-) or ionotropic nicotinic (N-type) receptors.

(a). Muscarinic receptors: Four sub types of muscarinic and one nicotinic receptor have been found in on cerebral arteries. Using isolated vessel preparations, Furchgott and Zawadzki (1980) demonstrated that the vasodilatory effect of ACh is endothelium-dependent. ACh acts on muscarinic receptors expressed on endothelial cells that stimulate endothelial NO-synthase activity(eNOS), releasing NO, which then triggers vasodilatation (Figure 6; (Wahl & Schilling, 1993)).

The endogenous origin of ACh for this mechanism is unclear, as it is rapidly broken down by AChE in the blood. Release from perivascular nerves is a possibility, as tritiated ACh applied to the outside of intact vessels readily crossed the media and appeared in the luminal perfusion fluid (Gonzalez *et al.* (1997). An endothelial origin has also been suggested, as endothelial cells contain ChAT and can produce and release ACh (Parnavelas *et al.*, 1985; Kawashima *et al.*, 1990), but what controls this is unclear.

-Insert Table 1 here-

(ACh receptors)

It is likely that muscarinic M_3 receptors located on endothelial cells mediate the NO induced vasodilatation, at least in rabbits (Garcia-Villalon *et al.*, 1991), cats (Dauphin & Hamel, 1990) and human (Dauphin & Hamel, 1992). In the absence of endothelium, ACh (exogenously applied or released by transmural stimulation) induces vasoconstriction via M_1 receptors located on the vascular smooth muscle cells (VSM). M_1 receptors have been identified in VSM of cerebral arteries including basilar arteries of rats (Phillips *et al.*, 1997) and in numerous other species including cat (Dauphin *et al.*, 1991), pig (Garcia-Villalon *et al.*, 1991) and human (Dauphin & Hamel, 1992). Since M_1 receptors have a lower affinity for ACh than M_3 receptors, higher concentrations of ACh are required for their activation and this may provide a mechanism for the differential vasomotor actions of ACh. From a study of pial vessels in mice (Shimizu *et al.*, 1993), there is evidence that the M_1 receptor mechanism is uncoupled/non-functional in the presence of a functional endothelial M_3 receptor, explaining the apparent dominance of M_3 receptor induced vasodilation.

Pre-junctional M_2 receptors are located on nitroergic perivascular neurones and attenuate the vasodilation (or increase vasoconstriction) by inhibiting NO production and release. This

has been found in several species (pig, cat, monkey) and in various locations along the posterior (and anterior) cerebral arteries (van Charldorp & van Zwieten, 1989; Dauphin & Hamel, 1992; Phillips *et al.*, 1997; Toda *et al.*, 1997; Liu & Lee, 1999). In contrast, there is a total absence of M₂ receptors in human cerebral arteries (Dauphin & Hamel, 1992), explaining the absence of ACh induced contraction in man mentioned above.

The M₅ receptor was the last muscarinic receptor to be identified in cerebral arteries and as with M₁ and M₃ receptors it is post-synaptically located. The receptor was found on both smooth muscle cells and endothelium (Phillips *et al.*, 1997; Tayebati *et al.*, 2003). The mRNA content for M₅ in rat basilar artery was found to be equal to that of M₂ and M₃ receptors, and greater than M₁ receptor mRNA (Phillips *et al.*, 1997). Due to the lack of selective M₅ muscarinic receptor ligands, little is known of the functional significance of this receptor on cerebral arteries. To circumvent this problem, Yamada *et al.* (2001b) created a M₅ receptor knock out mouse, in which large cerebral arteries and arterioles did not dilate following topical application of acetylcholine (both in *in vivo* and in organ bath preparation). The effect was specific to the cerebral blood vessels, as peripheral vessels retained the ability to dilate. It has been suggested that some of the effects previously ascribed to the M₃ receptor may be mediated via the M₅ receptor (Elhusseiny & Hamel, 2000).

(b) *Nicotinic ACh receptors*: There is a dearth of evidence supporting a major role for nicotinic receptors on vertebrobasilar arteries. Though Dauphin and Hamel (1992) did not find nicotinic receptors in cat or human cerebral arteries, Edvinsson *et al.* (1977) found evidence of the α 7-NACh receptor on adrenergic nerves of cat middle cerebral arteries. As

its function is relevant to modulation of adrenergic signalling, it will be described in the 'cross talk' section (see below).

(iii) Vasoactive Intestinal Peptide. VIP has been found to co-localise with ChAT in some parasympathetic nerves and ganglia innervating the cerebral vasculature (Gibbins & Morris, 1988; Suzuki *et al.*, 1990a; Hardebo *et al.*, 1992; Uddman *et al.*, 1999) including the basilar artery (Miao & Lee, 1990; Saito *et al.* 1985). In basilar arteries from cats, only 5% co-localisation between VIP and ChAT positive fibres was found despite these running closely to each other (Miao & Lee, 1990). The density of VIP innervation of cerebral arteries is species dependent with highest densities in the cat, rat and rabbit (Duckles & Said, 1982; Gibbins *et al.*, 1984b; Zhang *et al.*, 1991) and relatively sparse in human (Edvinsson *et al.*, 1987; Edvinsson *et al.*, 1994b). This confers with a study by White (1987) in which 87% of human basilar arteries pre-contracted with prostaglandin $F_{2\alpha}$ ($PGF_{2\alpha}$) displayed a sustained concentration dependent vasodilatation when exposed to VIP. Despite the lack of VIP positive fibre staining on vertebral and basilar arteries in dogs (Edvinsson *et al.*, 1980), the vascular conductance of vertebral arteries was increased by 100% on exposure to VIP (Blitz & Charbon, 1983). These data emphasise the importance of this peptide in mediating vasodilatation of vertebrobasilar arteries but the neural source remains to be ratified.

There are two VIP receptors with similar affinity for VIP: VIP₁ (VPAC₁) and VIP₂ (VPAC₂). In addition to these, VIP can also act via the natriuretic peptide clearance receptor (NPR-C) (Akiho *et al.*, 1995; Murthy *et al.*, 2000; Grant *et al.*, 2005). Immunohistochemical labelling for VPAC₁ in rat cerebral arteries found it on the surface of smooth muscle cells in the medial layer only (Fahrenkrug *et al.*, 2000). This was confirmed using *in situ* hybridization

mRNA, where VPAC₁ and VPAC₂ located into smooth muscles cells in posterior (and anterior) cerebral vessels (Baun *et al.*, 2011). Similar results have been obtained in humans, using RT-PCR against the two VPAC receptors (Knutsson & Edvinsson, 2002). In pig basilar arteries, mRNA is present for VPAC₁, VPAC₂ and NPR-C. Using immunocytochemistry, VPAC₁ was found in the endothelium, VPAC₂ on the outer layers of SMCs and endothelium and NPR-C throughout the artery including the perivascular nerves themselves (Grant *et al.*, 2005).

The vasodilatory effect of VIP may or may not involve a NO dependent mechanism.

Denuded cerebral arteries from sheep and cow showed blunted VIP induced vasodilation, indicating the presence of a mechanism involving NO of endothelial origin (Gaw *et al.*, 1991; Gonzalez *et al.*, 1997). VIP stimulation of isolated rat basilar arteries results in vasodilation, which is inhibited by the nNOS selective antagonist: 7-nitroindazole (7-NI), indicating involvement NO from perivascular nitrergic nerves (Seebeck *et al.*, 2002). VIP receptors present on SMC, however, induce vasodilation directly, via an NO independent mechanism, probably through the cAMP pathway (Grant *et al.*, 2005).

(iv) Pituitary adenylate cyclase-activating peptide (PACAP) belongs to the same family of glucagon/secretin neuropeptides as VIP and shares 68% amino acid sequence homology.

The two co-transmitters are expressed in different subpopulations of parasympathetic neurones. In addition, PACAP is also expressed by both trigeminal sensory neurones and sympathetic neurons making its association with the parasympathetic nervous system equivocal (Sundler *et al.*, 1996). PACAP mRNA is expressed in most otic and PPG cells in the rat (Sundler *et al.*, 1996). PACAP acts through 3 different G protein coupled receptors: the VIP receptors: VPAC₁ & VPAC₂, and PAC₁, all increase adenylate cyclase activity. There is

evidence of vasodilatory effects of PACAP in rat basilar arteries mediated by nitregic nerves (Seebeck *et al.*, 2002). Although PACAP is up to 100x more potent as a vasodilator than VIP (Kashimoto *et al.*, 1996), PACAP fibre innervation of cerebral arteries is less dense than that of VIP and NO (Warren *et al.*, 1991; Edvinsson *et al.*, 2001). In pre-constricted rat basilar arteries a stronger vasodilatory capacity of VIP than PACAP was found, whereas VIP and PACAP acted similarly in the anterior vessel segments, which may indicate a differential receptor distribution between these arteries (Baun *et al.*, 2011).

(v) Peptide histidine isoleucine. VIP-like neuropeptides such as peptide histidine isoleucine (PHI) in rat and the human equivalent peptide histidine methionine (Linder *et al.*, 1987) are also coexpressed in the VIP population of parasympathetic fibres on basilar arteries and have weaker vasodilatory properties relative to VIP (Suzuki *et al.*, 1984; Uddman *et al.*, 1993).

Cross talk between parasympathetic and sympathetic nerves

Parasympathetic and sympathetic nerves are often intertwined on cerebral arteries (Owman *et al.*, 1974) or run in parallel within the same perineural sheath (Iwayama *et al.*, 1970; Edvinsson *et al.*, 1972a). Axo-axonal contact distances between sympathetic and parasympathetic fibres can be as little as 25 nm, compared to the 100 nm seen for neuro-muscular contacts in the same vessels (Edvinsson *et al.*, 1972a). This close apposition suggests cross talk between parasympathetic and sympathetic nerve fibres.

-Insert Figure 5 here-

(Close apposition of NA and Ach nerves)

Support for interactions between the parasympathetic and sympathetic innervations to cerebral arteries is that they develop simultaneously in both mice and rats (Kobayashi *et al.*, 1981; Tsai *et al.*, 1992) and human: (Kawamura & Takebayashi, 1994). In pre- and postnatal rats, parasympathetic and sympathetic fibres (visualised with VIP and NPY markers, respectively) show similar patterns of development in their distribution, density and pattern (Tsai *et al.*, 1992). This close association in development was not observed for sensory fibres (visualised using CGRP), their development occurs after sympathetic innervation (Tsai *et al.*, 1989). There is not the same degree of close apposition between sensory fibres with either sympathetic or parasympathetic nerves (Tsai *et al.*, 1989; Edvinsson *et al.*, 1994a).

Functional studies in basilar arteries from rat and pig have found that NA and ACh have limited vasoreactivity, yet adrenergic nerves express nicotinic α -7 NACH receptors, which upon stimulation, cause increases in intracellular calcium leading to noradrenaline release from their varicosities (Si & Lee, 2001; Si & Lee, 2002). This triggers vasodilatation via activation of β_2 -adrenergic receptors; vasoconstriction does not occur as there is paucity of α -adrenoceptors. β_2 -adrenoceptors are located on the intertwined acetylcholinergic-nitrergic fibres leading to NOS activation and NO production and stimulation of guanylyl cyclase (Fig 7). Removal of the endothelium does not affect the arteries ability to dilate, indicating the source of NO must be neuronal (Lee, 1980; Toda, 1982). Indeed, NO is stored and released with ACh. ACh acts on nitrergic neurones stimulating prejunctional M_2 receptors and inhibits NO synthesis, thus modulating NO release (Liu *et al.*, 2002). Following unilateral excision of the superior cervical ganglion in rats, NO mediated vasodilation in cerebral arteries was reduced, indicating a dependence on an intact sympathetic innervation (Smith *et al.*, 2002).

-Insert Figure 6 here-

(Mechanism of crosstalk)

Cross talk may have a role in the development and maintenance of autonomic fibres and their phenotype. For example, after sympathectomy: (i) parasympathetic fibres proliferate and make more direct contacts with vascular smooth muscle cells, whereas prior to sympathectomy they mainly contact sympathetic fibres (Smith *et al.*, 2002); (ii) parasympathetic cells can display increased dopamine- β -hydroxylase (DBH) immunoreactivity (though this is functionally irrelevant), and this is matched by a similar decrease in the number of PPG neurones expressing parasympathetic markers (Mione *et al.*, 1991; Fan & Smith, 1993); (iii) both NOS and VIP activity in the PPG is decreased, resulting in a reduction in NOS dependent vasodilation in various target areas (Fan & Smith, 1993; Smith *et al.*, 2002).

In face of increased sympathetic drive as occurring during “fight or flight” situations vertebrobasilar arteries may thus vasodilate (instead of constrict) via the mechanisms described above to ensure the brainstem and or cortex remains well perfused at times when peripheral organs experience vasoconstriction and reduced vascular conductance (Lee *et al.*, 2011).

Other proposed functions of parasympathetic innervation to cerebral arteries *in vivo*

With such a dense parasympathetic innervation of the cerebral arteries it is a natural question to ask: What are the physiological circumstances in which these nerves are activated?? Beside the suggested function in ensuring adequate CBF during periods of high sympathetic activity, what else does it do? The question has not been answered equivocally,

though various clues to its function are appearing: As mentioned earlier activation of the parasympathetic innervation causes vasodilation. Thus it is likely to be involved in any process requiring centrally regulated increases in CBF that are independent of the endothelial NO system. Examples include, increased brain metabolism, thermoregulation (it is essential to keep the brain thermoregulated) and hypercapnia.

Electrical stimulation of the PPG in rats increases cerebral blood flow by 40-50% on the ipsilateral side as measured by Doppler flow over the parietal cortex and tissue PO₂ (Seylaz *et al.*, 1988; Talman *et al.*, 2007). In dogs, stimulation of the PPG and the superficial petrosal evoked vasodilatation in the middle cerebral artery, posterior communicating arteries and basilar artery (D'Alecy & Rose, 1977; Toda *et al.*, 2009). Anti-cholinergic drugs fails to attenuate the response, emphasising that the vasodilatory response is mediated via a non-cholinergic mechanism (Nakai *et al.*, 1993).

What evidence is there for vasodilatory tone? Following removal of the PPG in the awake rat Suzuki *et al.* (1990b) reported no change in blood flow, suggesting an absence of tone. However, these data were not confirmed by Boysen *et al.* (2009), who saw ablation of the PPG evoking a 30% decrease in blood flow increasing cerebrovascular resistance (CVR) in rats. Given that the muscarinic antagonist, atropine, does not affect cerebral blood flow, non-cholinergic transmission must underlie this tone (Toda *et al.*, 1993; Yoshida *et al.*, 1993; Toda *et al.*, 2000a). Further evidence of tonic parasympathetic tone in the form of reduced blood flow upon NO inhibition has also been observed in monkeys (Toda *et al.*, 2000b) and dogs (Toda *et al.*, 1993; Yoshida *et al.*, 1993; Toda *et al.*, 2000a), where the vasodilatory responses to nerve stimulation was found to be attenuated by the NO inhibitor N^G-nitro-L-arginine (L-NA). Indeed, NO of neuronal origin has been associated with vasodilation and

decreased cerebral vascular resistance (Toda & Okamura, 2003). In nNOS deficient mice a reduced vasodilator response following stimulation of both efferent and afferent nerves was observed, an effect not seen in eNOS or iNOS deficient mice (Toda *et al.*, 2009). Thus, there appears to be an apparent loss of ACh dependency as primary transmitter in the parasympathetic innervation and transfer to NO mediated vasodilatory processes, which is distinct to the innervation of peripheral arteries.

Parasympathetic innervation from the PPG acting directly or indirectly on nociceptive sensory fibres have also been suggested to participate in regulation of blood brain barrier (BBB) permeability (Delépine & Aubineau, 1997; Yarnitsky *et al.*, 2004a; Yarnitsky *et al.*, 2004b) and development of cluster headaches/migraine (Yarnitsky *et al.*, 2003). Additionally, it is involved in the dilation of cerebral arteries following “breakthrough of autoregulation” in acute hypertension (Agassandian *et al.*, 2003).

Parasympathetic targets with clinical relevance for the cerebral vasculature

The critical importance of the brainstem for life means that its perfusion needs to be tightly regulated. At rest, autonomic tone of the cerebral vessels is small (Heistad & Marcus, 1978; Heistad *et al.*, 1978). However, it is reasonable to assume that any imbalance in the activity of autonomic nerves innervating the posterior cerebral circulation could have pathological implications for blood flow control and be a potential clinical target for various conditions in which cerebral blood flow is impeded, such as stroke or hypertension. The importance of the brainstem is highlighted by the finding that during extreme acute hypotension, cerebral blood flow is prioritised to the brainstem (Mueller *et al.*, 1977). In this regard, we consider how disease is affected by the parasympathetic regulation of the cerebral vasculature and

ways in which modulation of parasympathetic input to the cerebral arteries may be used to alleviate pathology.

During pathological increases of blood pressure above the upper limit of autoregulation, cerebral vessels dilate. Denervation of parasympathetic fibres originating from the PPG dampens this effect suggesting a baroreceptor mediated cerebral vasodilatation in rats (Talman & Dragon, 2000). It was later found that neuronal NOS (nNOS) drove the observed dilatory response; using a specific nNOS inhibitor the breakthrough dilatation that occurs above the autoregulatory threshold was attenuated (Talman *et al.*, 2007). One possibility is that the shift in the autoregulatory curve to the right in hypertension (e.g. as in the SHR) may reflect reduced parasympathetic nitrenergic input.

The importance of NO in cerebral vascular regulation is highlighted by the association of the endogenous NOS inhibitor - asymmetric dimethylarginin (ADMA) which is elevated in conditions of hypoperfusion such as stroke and hypertension (Yoo & Lee, 2001; Napoli *et al.*, 2004; Kielstein & Zoccali, 2005). By targeting ADMA availability (Fan *et al.*, 2013) and reducing plasma levels (Fan *et al.*, 2013; Tsai *et al.*, 2014) hypertension is prevented in young SHRs. Furthermore, ADMA infusion at levels that do not affect systemic blood pressure decreased CBF and induces vascular stiffness in healthy volunteers (Kielstein *et al.*, 2006). In rats topical application of ADMA results in basilar vasoconstriction and an inhibition of ACh mediated vasodilatation and in rabbits it has the same effect in cortical vessels (Faraci *et al.*, 1995). Thus, NO in conjunction with parasympathetic innervation plays a major role in maintaining blood flow to the brain in health and disease states. The role of ADMA in ischemic stroke and its potential as a predictive disease marker has been discussed and reviewed elsewhere (Chen *et al.*, 2012).

The endogenous NOS inhibitor monomethylarginine (L-NMMA), has been reported to be present at lower concentrations than ADMA in the circulation (Vallance *et al.*, 1992). Like ADMA, L-NMMA was shown to reduce cortical blood flow when injected into the internal carotid artery without affecting systemic blood pressure (Thompson *et al.*, 1996). When challenged with hypercapnia, the chemoregulatory response normally resulting in increased cerebral blood flow in monkeys, was prevented by prior intracarotid injection of L-NMMA. The same was not true for cerebral autoregulation due to changes in blood pressure (Thompson *et al.*, 1996). This result provides further evidence for the role of NO in control of cerebral blood flow.

Nitric Oxide is synthesised by NOS from L-arginine. The active site of NOS is subject to inhibition by the asymmetric methylarginines, ADMA and L-NMMA. These inhibitors could be a good clinical target in conditions where the synthesis or bioavailability of NO is compromised (Leiper & Nandi, 2011). However, it is not specific to nNOS but also work on eNOS.

The finding of markers associated with cardiovascular risk, and possibly autonomic imbalance, such as ADMA (Faraci *et al.*, 1995), could provide other therapeutic pathways with targeted prophylactic medicine. In humans, sulfhydryl ACE inhibitor to reduce blood pressure also restored the NO/ADMA balance in hypertension (Napoli *et al.*, 2004). Similarly, in SHR 10 week treatment with an ACE-inhibitor not only decreased blood pressure but improved the vasodilatory response of the basilar artery and blood flow during hypotension as well as the NO-mediated dilatation during topical application of ACh; these data all suggest an improvement in endothelial function of the basilar artery (Toyoda *et al.*,

1998). Furthermore, NO or NO donor administration in stroke models has been shown to reduce lesion size and increase cerebral perfusion (Willmot *et al.*, 2005).

The direct effects of parasympathetic innervation can be seen during a hypotensive challenge in SHR in which forebrain blood flow was reduced after chronic parasympathetic denervation compared to sham controls (Koketsu *et al.*, 1992). These findings were not corroborated in normotensive Sprague Dawley rats (Branston *et al.*, 1995). The parasympathetic system is involved in vasodilatation at the lower end of autoregulation (Branston *et al.*, 1995). To what extent this is true for the posterior circulation of the brainstem remains to be established. In stroke, parasympathetic denervation exacerbates infarct size following middle cerebral artery occlusion (Kano *et al.*, 1991; Koketsu *et al.*, 1992). This evidence supports a role for parasympathetic nerves in protecting against brain ischemia. In this context, stimulation of the PPG provided protection by increasing CBF and reducing the infarct size following cerebral artery occlusion (Henninger & Fisher, 2007). A reduction in ischemic area as well as protection of blood-brain barrier integrity were found in a model of photothrombotic stroke with PPG stimulation (Levi *et al.*, 2012).

It is worth considering the potential clinical importance of cross talk between sympathetic and parasympathetic branches mentioned above, as it can result in alterations in autonomic regulation in pathological circumstances. For instance stimulation of sympathetic nerves in normotensive rats results in basilar artery vasodilatation and increased blood flow, the same is not true for hypertensive animals or animals with chronic superior cervical ganglion denervation (Chang *et al.*, 2012). The attenuated response is likely to be, at least in part, due to a reduction in the sympathetic to parasympathetic terminal density at the level of

the basilar artery in the hypertensive animals (Lee & Saito, 1986) or a lack of sympathetic input (Chang *et al.*, 2012). Conversely, an overactive sympathetic and underactive parasympathetic nervous system limits basilar artery dilatation in face of a hypotensive challenge (Barry *et al.*, 1982). This may cause ischemic damage especially in conditions where cerebrovascular resistance is high (e.g. SHRs), (Paton *et al.*, 2009; Cates *et al.*, 2011; Cates *et al.*, 2012). Furthermore the limited vasodilatation of the basilar artery (Toyoda *et al.*, 1998) contributes to the shift of the autoregulatory curve to higher BP levels in hypertension (Barry *et al.*, 1982) further increasing the susceptibility of hypoperfusion.

In case of ischemic stroke, stimulation of the parasympathetic nervous system to increase cerebral blood flow and restrict ischemia has clinical potential particularly considering the limiting treatments currently available and the restrictions of their use, as is the case with thrombolysis therapy (Cheyuo *et al.*, 2011). Stroke in the vertebrobasilar circulation is a clinical emergency and accounts for about 20% of all ischemic strokes (Schoen *et al.*, 2011) and has a high morbidity and mortality rate. Evidence shows that parasympathetic input provides some protection from ischemic damage (Kano *et al.*, 1991; Koketsu *et al.*, 1992; Henninger & Fisher, 2007; Cheyuo *et al.*, 2011; Levi *et al.*, 2012). Importantly this can be effective even after 24 hours of stroke onset (Solberg *et al.*, 2008; Levi *et al.*, 2012). The mechanisms of action require further investigation, as does the relevance of this treatment to the vertebrobasilar circulation. A comprehensive review on the potential of harnessing the parasympathetic system as a therapy in stroke has been published recently (Cheyuo *et al.*, 2011). Whether this can also be targeted to assist with vertebrobasilar artery hypoperfusion in neurogenic hypertension (Cates *et al.*, 2012; Marina *et al.*, 2015), for example, remains unknown but holds therapeutic potential. Indeed, the Brainsgate Ischemic

Stroke System comprises an stimulator implanted adjacent to the PPG via the greater palatine canal under local anaesthesia (Khurana *et al.*, 2009). This device is currently undergoing phase 3 trials (Clinical Trials.gov: NCT00826059 & NCT01874093). Another device directed to stimulation of the otic ganglion has also been described (Shalev & Gross, 2010). By extrapolation, once the source(s) of input to the posterior cerebral circulation has been clarified, these nerves/ganglia could be targeted in a similar fashion.

Systemically, activation of the parasympathetic efferent vagus nerve has been shown to inhibit pro-inflammatory cytokine release and decrease the inflammatory response. This function is coined “the cholinergic anti-inflammatory pathway” (Borovikova *et al.*, 2000; Pavlov *et al.*, 2003; Tracey, 2007). Evidence suggest it works via nicotinic $\alpha 7$ -acetyl cholinergic receptor stimulation (Wang *et al.*, 2003) and has been reported to have beneficial effects in a plethora of diseases from irritable bowel syndrome to depression (Wang *et al.*, 2004; Rush *et al.*, 2005; van Westerloo *et al.*, 2006; Ghia *et al.*, 2007; van Maanen *et al.*, 2009) as well as improving kidney transplant outcomes (Hoeger *et al.*, 2010). We propose that activation of the parasympathetic pathways innervating the cerebral arteries could also yield a similar anti-inflammatory response. In the anterior circulation this may be relevant for Alzheimer’s disease (de la Torre, 2002; Reale *et al.*, 2004) and for posterior cerebral arteries: hypertension, which has been linked to inflammatory activity in brainstem walls (Waki *et al.*, 2007; Waki *et al.*, 2010; Zubcevic *et al.*, 2011; Waki *et al.*, 2013).

In this regard, both VIP and PACAP have been reported to exert similar anti-inflammatory effects through the VPAC1 receptor, which could be a relevant mechanism for the

parasympathetic innervation of cerebral arteries (Leceta *et al.*, 2000; Martinez *et al.*, 2002; Grimm *et al.*, 2003).

Yarnitsky *et al.* (2005) have shown that electrical stimulation of the PPG can prevent cerebral vascular spasm after subarachnoid haemorrhage and improve blood flow.

Acetylcholine (given via intracisternal injection) can counter effects of cerebral vasospasm in basilar arteries after subarachnoid haemorrhage by suppressing the production of cytokines (Song *et al.*, 2014). However, it is worth noting that PPG stimulation has also been reported to disrupt the BBB and induce neurogenic inflammation via a reflex involving nociceptive sensory fibres (Delépine & Aubineau, 1997). A similar mechanism has also been ascribed to pain in migraine: Yarnitsky *et al.* (2003) reported that PPG activation during migraine is involved in sensitization of intracranial nociceptors by direct or indirect activation of sensory pain fibres by the release of ACh, VIP and NO. This effect of PPG stimulation has been used clinically to deliver macromolecules through the BBB (Yarnitsky *et al.*, 2004a). Hence the (over-) stimulation of the parasympathetic innervation to the cerebral arteries may not only yield positive effects. Clinical application will require careful adaptation depending on the effect sought.

Finally, and hypothetically, peptide targeting of the cerebral vasculature may also be possible. Using phage display proteins were identified that targeted the endothelial cells of the aorta and heart (Greig *et al.*, 2010). Using such a technique may allow selective targeting of the cerebral arteries and may even differentiate between posterior versus the anterior circulation. Such targeting might allow enhancement of efficacy of vasodilating transmitter substances, for example.

Concluding remarks

Control of blood flow to the brain is highly specialised and regionally specific. Uniquely, resistance is controlled in part by the main supply vessels as well as the microcirculation. We have discussed how the Circle of Willis and the vertebrobasilar circulations have different patterns of nervous innervation from parasympathetic, sympathetic and sensory systems. Because of similar phenotypes, we stress caution as to how to separate parasympathetic from sensory innervations immunocytochemically. Whilst the parasympathetic innervation of the anterior cerebral circulation is well described, this review highlights the need for clarification of the sources of parasympathetic innervation to the posterior cerebral circulation. It is clear that there are multiple sources of parasympathetic innervation and these change with age and in disease states and between species. The distinct sources of origin provides an opportunity for targeted regulation affecting sub-components of the cerebral circulation, which may be relevant to the treatment of stroke. The review highlights the number of transmitters systems used by the parasympathetic nerves innervating the cerebral vessels (e.g. ACh, NO, VIP, PACAP & PHI); this may provide fail safe mechanisms, state-dependent control, and regulation of the magnitude of the vasoactive response, for example. The evidence is unequivocal that there is a strong interdependence between the sympathetic and parasympathetic innervation both developmentally and functionally as they modulate each other. An example is the basilar artery vasodilatation induced by sympathetic nervous stimulation triggered by NO release from parasympathetic terminals. Malfunction of one of these systems will functionally alter the performance of the other. Thus, therapy targeting the parasympathetic nervous system to increase brain blood flow should not only be assured of a functional parasympathetic but also sensory and sympathetic innervation.

Innovative methods are being used to selectively stimulate specific nerves and ganglia innervating components of the cerebral circulation to increase blood flow. Whether specific endothelial markers on aspects of the cerebral circulation can be identified and target to modulate parasympathetic mediated vasodilatation remains a project for future scientific enquiry.

Additional information

Competing interests

The authors have no competing interests to report.

Author contributions

Eva v.L. Roloff: Research, 1st draft of article, contribution to figures and revision; Ana Tomiak-Baquero: Figures, research and contribution to article (sections in 1st draft), revision; Sergey Kasparov: Contributions to concept, grantholder & revision, Julian F.R. Paton: Developer of concept, contribution to main article and major revision, and primary grantholder. All authors approve the final version of the manuscript.

Funding

This article was written with the support of The British Heart Foundation (BHF) RG/12/6/29670

Author profiles:

EvLR currently is a research associate with speciality in systems physiology and neuroscience. She holds a Masters in Biology from Copenhagen, DK (1995) and a PhD from Aberdeen, Scotland (2003). **AMTB** completed her PhD in 2013 at the University of Bristol. She is currently a Research Associate working on brain blood flow control in rodents with hypertension. **SK** holds a MD PhD (Moscow State University, Russia) and has a Chair in Molecular Neuroscience. Current interests are in astrocyte signalling. **JFRP** completed his PhD in 1987 (University of London) and trained in London, USA and Germany. In 2004 he was awarded a British Heart Foundation Lectureship. He leads a translational research group in cardiovascular and respiratory disease.

Figure Legends

Figure 1. Human angiogram and schematic of cerebral circulation.

(A) CT scan showing the anatomy of the cerebral arterial supply in humans. The image was kindly provided by Dr Nathan Manghat from University Hospital Bristol Trust. (B) Labelled schematic representation of (A) including origins of the circulation form the aortic arch, Figure modified from Schuenke *et al.* (2010). Abbreviations: A, anterior; R, right; P, posterior; L, left.

Figure 2. Species variation in cerebral artery architecture.

Schematic representation of the cerebral arterial tree and difference in contribution to anterior and posterior circulation by the ECA and ICA in (A) rats, (B) goat/sheep and (C) rabbit. Notice the presence of the carotid rete in (B) goat/sheep (red arrow), the lack of ICA and the existence of a V-Oa in the same species. CoW and VA labelled for orientation purposes. Abbreviations: APAr, ascending pharyngeal artery residual; BA, basilar artery; CCA, common carotid artery; CoW, Circle of Willis; SA, spinal artery; VA, vertebral artery; V-Oa, vertebral-occipital anastomoses. Figure adapted from Daniel *et al.* (1953), Andersson and Jewell (1956) and Baldwin and Bell (1963).

Figure 3. Approximate spatial contribution to cerebral blood supply by carotid and vertebral arteries in different species.

Schematic representation of the cerebral blood supply contribution by the carotid and vertebral arteries in (A) man, (B) rat, (C) goat/sheep, (D) rabbit, (E) dog, (F) cat and (G) calf. Image based and adapted from Baldwin and Bell (1963) and Bralet *et al.* (1977).

Figure 4. Parasympathetic innervation to cerebral vasculature.

(A) Schematic overview of the anatomic position of various parasympathetic ganglia which are sources of parasympathetic input to the cerebral arteries. The level of parasympathetic innervation to the cerebral arteries varies as shown by (B) VIP innervations in the posterior communicating artery (dense innervation) and basilar artery (moderate innervation); image from Hara *et al.* (1985); x172 in original article, reproduced with permission. (C) Schematic representation of parasympathetic innervations to the rostral and posterior cerebral vasculature. The gradient represents reported density of innervation, checkered filling indicates unreported innervation. Note that species variation regarding innervation is not depicted in the figure. CmG, Carotid miniGanglia; CS, Cavernous Sinus; OG, Otic Ganglia; PTG; Pterygopalatine Ganglia; VIP, vasoactive intestinal peptide. Original figure collated from data from Gibbins *et al.* (1984a); Hara *et al.* (1985); Keller *et al.* (1985); Shimizu (1994); Kadota *et al.* (1996); Bleys *et al.* (2001); Ayajiki *et al.* (2012).

Figure 5. Electron microscopy image showing the close proximity of sympathetic and parasympathetic terminals on a cerebral artery.

Image of the anterior cerebral artery of cat showing the close apposition of a parasympathetic (Ch, cholinergic) and sympathetic (A, adrenergic) neurons. Schwann cell is visible at the bottom. Image from Edvinsson *et al.* (1972b). x60000 in original reference, with permission.

Figure 6. Mechanisms of endothelial dependent cross-talk leading to NO induced vasodilatation.

There are multiple sources of nitric oxide production including the endothelium. (A) ACh exerts endothelium dependent vasodilatation acting on muscarinic (M) type 3 and/or 5 receptors on endothelial cells inducing an increase in intracellular calcium which leads to NO production via eNOS activity. ACh still evokes a dilatation in denuded vessels suggesting a non-endothelial source of nitric oxide (NO). (B) Norepinephrine (NE) can stimulate the production of NO from cholinergic/nitrergic neurons via β_2 receptors leading to increases in Ca^{2+} and calcium channel stimulation. The NO diffuses to nearby smooth muscle cells causing relaxation. Modulation of NO release in this system is achieved by ACh mediated NO inhibition via pre-junctional muscarinic (M) type 2 receptors on nitrergic neurons.

Abbreviations: ACh, acetylcholine; cGMP, cyclic guanosine monophosphate; GC, guanylate cyclase; GTP, guanosine triphosphate; NO, nitric oxide; NOS, nitric oxide synthase. Figure based and adapted from Wahl and Schilling (1993), Zhang *et al.* (1998b), Liu *et al.* (2000) and Si and Lee (2002).

Tables

Receptor subtype	Cellular location	Vessel location	Effect	Proportion of total receptors	Species	Reference
M ₁ -AChR	SMC	MCA	Vasoconstriction	15-27%	Cat	(Dauphin <i>et al.</i> , 1991)
				26-54%	Human	(Dauphin & Hamel, 1992)
M ₂ - AChR	Perivascular nerves	Absent!! BA	Auto regulatory	35% 0%	Cat & human	(Dauphin & Hamel, 1992)
	Nitrgenic nerves		NO-modulatory-inhibition of NO - vasoconstriction	-	Pig	(van Charldorp & van Zwieten, 1989; Liu & Lee, 1999)
	Pre-junctionally NO-nerves	MCA ⁴	Attenuate vasodilation by inhibiting NO-release from nitrgenic nerves		Monkey	(Toda <i>et al.</i> , 1997)
M ₃ - AChR	Endothlial	MCA	Vasodilation	10-18%	Cat & human	(Dauphin & Hamel, 1990; Dauphin <i>et al.</i> , 1991)
M ₅ - AChR	?	BA	?	?	Rat	(Phillips <i>et al.</i> , 1997; Tayebati <i>et al.</i> , 2003)
	Endothelium & SMC (less) media	CoWs	Vasodilation		GM-mouse	(Yamada <i>et al.</i> , 2001a)
α ₇ -N AChR				none	Cat & human	(Dauphin & Hamel, 1992)
		BA	Modulation of sympathetic		Pig	(Zhang <i>et al.</i> , 1998a; Si & Lee, 2001)
		BA	perivascular neurones		Rat	(Chang <i>et al.</i> , 2012)
α ₃ β ₂ - N AChR			Modulation of sympathetic perivascular neurones		Rat	(Lee <i>et al.</i> , 2011)

Table 1. Cholinergic receptors identified on cerebral arteries.

⁴Endothelial denuded vessel prep

REFERENCES

- Agassandian K, Fazan VS, Margaryan N, Dragon D, Riley J & Talman W. (2003). A Novel Central Pathway Links Arterial Baroreceptors and Pontine Parasympathetic Neurons in Cerebrovascular Control. *Cellular and Molecular Neurobiology* **23**, 463-478.
- Akiho H, Chijiwa Y, Okabe H, Harada N & Nawata H. (1995). Interaction between atrial natriuretic peptide and vasoactive intestinal peptide in guinea pig cecal smooth muscle. *Gastroenterology* **109**, 1105-1112.
- Alpers BJ, Berry RG & Paddison RM. (1959). Anatomical studies of the circle of Willis in normal brain. *AMA archives of neurology and psychiatry* **81**, 409-418.
- Andersson B & Jewell P. (1956). The distribution of carotid and vertebral blood in the brain and spinal cord of the goat. *Quarterly journal of experimental physiology and cognate medical sciences* **41**, 462-474.
- Ando K. (1988). Distribution and origin of vasoactive intestinal polypeptide (VIP)-immunoreactive, acetylcholinesterase-positive and adrenergic nerves of the cerebral arteries in the bent-winged bat (Mammalia: Chiroptera). *Cell and tissue research* **251**, 345-351.
- Arbab MAR, Wiklund L & Svendgaard NA. (1986). Origin and distribution of cerebral vascular innervation from superior cervical, trigeminal and spinal ganglia investigated with retrograde and anterograde WGA-HRP tracing in the rat. *Neuroscience* **19**, 695-708.
- Ayajiki K, Kobuchi S, Tawa M & Okamura T. (2012). Nitroergic nerves derived from the pterygopalatine ganglion innervate arteries irrigating the cerebrum but not the cerebellum and brain stem in monkeys. *Hypertension research* **35**, 88-92.
- Baker MA. (1982). Brain cooling in endotherms in heat and exercise. *Annual review of physiology* **44**, 85-85.
- Baldwin B & Bell F. (1963). The anatomy of the cerebral circulation of the sheep and ox. The dynamic distribution of the blood supplied by the carotid and vertebral arteries to cranial regions. *Journal of anatomy* **97**, 203.
- Barnett CH & Marsden CD. (1961). Functions of the mammalian carotid rete mirabile. *Nature* **191**, 88-89.

- Barry DI, Strandgaard S, Graham DI, Braendstrup O, Svendsen UG, Vorstrup S, Hemmingsen R & Bolwig TG. (1982). Cerebral blood flow in rats with renal and spontaneous hypertension: Resetting of the lower limit of autoregulation. *Journal of cerebral blood flow and metabolism* **2**, 347-353.
- Baumbach GL & Heistad DD. (1983). Effects of sympathetic stimulation and changes in arterial pressure on segmental resistance of cerebral vessels in rabbits and cats. *Circulation research* **52**, 527-533.
- Baun M, Hay-Schmidt A, Edvinsson L, Olesen J & Jansen-Olesen I. (2011). Pharmacological characterization and expression of VIP and PACAP receptors in isolated cranial arteries of the rat. *European journal of pharmacology* **670**, 186-194.
- Beausang-Linder M & Bill A. (1981). Cerebral circulation in acute arterial hypertension—protective effects of sympathetic nervous activity. *Acta physiologica Scandinavica* **111**, 193-199.
- Bevan JA, Bevan RD, Florence VM, McCalden TA, Pearce WJ, Rowman RA & Winkquist RJ. (1981). Specialization in cerebral and extracerebral neurovascular mechanisms. *FASEB J* **40**, 2301-2305.
- Bevan JA, Moscovitz M, Said SI & Buga G. (1984). Evidence that vasoactive intestinal polypeptide is a dilator transmitter to some cerebral and extracerebral cranial arteries. *Peptides* **5**, 385-388.
- Bill A & Linder J. (1976). Sympathetic control of cerebral blood flow in acute arterial hypertension. *Acta physiologica Scandinavica* **96**, 114-121.
- Bleys RL, Groen GJ & Hommersom RF. (1996a). Neural connections in and around the cavernous sinus in rat, with special reference to cerebrovascular innervation. *The journal of comparative neurology* **369**, 277-291.
- Bleys RLAW & Cowen T. (2001). Innervation of cerebral blood vessels: Morphology, plasticity, age-related, and Alzheimer's disease-related neurodegeneration. *Microscopy research and technique* **53**, 106-118.
- Bleys RLAW, Cowen T, Groen GJ, Hillen B & Ibrahim NBN. (1996b). Perivascular nerves of the human basal cerebral arteries I. Topographical distribution. *Journal of cerebral blood flow and metabolism* **16**, 1034-1047.
- Bleys RLAW, Thrassivoulou C & Cowen T. (2001). Cavernous sinus ganglia are sources for parasympathetic innervation of cerebral arteries in rat. *Journal of cerebral blood flow and metabolism* **21**, 149-156.

- Blitz W & Charbon GA. (1983). Regional vascular influences of vasoactive intestinal polypeptide. *Scandinavian journal of gastroenterology* **18**, 755-763.
- Bogdan C. (2001). Nitric oxide and the immune response. *Nature immunology* **2**, 907-916.
- Borovikova LV, Ivanova S, Zhang M, Yang H, Botchkina GI, Watkins LR, Wang H, Abumrad N, Eaton JW & Tracey KJ. (2000). Vagus nerve stimulation attenuates the systemic inflammatory response to endotoxin. *Nature* **405**, 458-462.
- Boysen NC, Dragon DN & Talman WT. (2009). Parasympathetic tonic dilatory influences on cerebral vessels. *Autonomic neuroscience* **147**, 101-104.
- Bralet A, Rochette L & Bralet J. (1977). [Evaluation of the carotid blood distribution in the rat using radioactive microspheres (author's transl)]. *Experientia* **33**, 350-352.
- Branston NM, Umemura A & Koshy A. (1995). Contribution of cerebrovascular parasympathetic and sensory innervation to the short-term control of blood flow in rat cerebral cortex. *Journal of cerebral blood flow and metabolism* **15**, 525-531.
- Bredt DS, Hwang PM & Snyder SH. (1990). Localization of nitric oxide synthase indicating a neural role for nitric oxide. *Nature* **347**, 768-770.
- Brown JO. (1966). The morphology of circulus arteriosus cerebri in rats. *The anatomical record* **156**, 99-106.
- Carvalho VC. (1985). Nerve cells in the human cavernous sinus. *Anatomischer Anzeiger* **159**, 29-32.
- Cates MJ, Dickinson CJ, Hart EC & Paton JF. (2012). Neurogenic hypertension and elevated vertebrobasilar arterial resistance: is there a causative link? *Current hypertension reports* **14**, 261-269.
- Cates MJ, Steed PW, Abdala AP, Langton PD & Paton JF. (2011). Elevated vertebrobasilar artery resistance in neonatal spontaneously hypertensive rats. *Journal of applied physiology* **111**, 149-156.
- Chang H-H, Lee Y-C, Chen M-F, Kuo J-S & Lee TJF. (2012). Sympathetic activation increases basilar arterial blood flow in normotensive but not hypertensive rats. *American journal of physiology - heart and circulatory physiology* **302**, H1123-H1130.
- Chen S, Li N, Deb-Chatterji M, Dong Q, Kielstein JT, Weissenborn K & Worthmann H. (2012). Asymmetric dimethyarginine as marker and mediator in ischemic stroke. *International journal of molecular sciences* **13**, 15983-16004.

- Cheyuo C, Jacob A, Wu R, Zhou M, Coppa GF & Wang P. (2011). The parasympathetic nervous system in the quest for stroke therapeutics. *Journal of cerebral blood flow & metabolism* **31**, 1187-1195.
- Chorobski J & Penfield W. (1932). Cerebral vasodilator nerves and their pathway from the medulla oblongata: With observations on the pial and intracerebral vascular plexus. *Archives of neurology & psychiatry* **28**, 1257-1289.
- D'Alecy LG & Feigl EO. (1972). Sympathetic control of cerebral blood flow in dogs. *Circulation research* **31**, 267-283.
- D'Alecy LG & Rose CJ. (1977). Parasympathetic cholinergic control of cerebral blood flow in dogs. *Circulation research* **41**, 324-331.
- Dahl E. (1973). The innervation of the cerebral arteries. *Journal of anatomy and physiology* **115**, 53-63.
- Daniel PM, Dawes JKD & Prichard MML. (1953). Studies of the carotid rete and its associated arteries. *Philosophical transactions of the Royal Society of London Series B, Biological sciences* **237**, 173-208.
- Dauphin F & Hamel E. (1990). Muscarinic receptor subtype mediating vasodilation in feline middle cerebral artery exhibits M3 pharmacology. *European journal of pharmacology* **178**, 203-213.
- Dauphin F & Hamel E. (1992). Identification of multiple muscarinic binding site subtypes in cat and human cerebral vasculature. *Journal of pharmacology and experimental therapeutics* **260**, 660-667.
- Dauphin F & Mackenzie ET. (1995). Cholinergic and vasoactive intestinal polypeptidergic innervation of the cerebral arteries. *Pharmacology & therapeutics* **67**, 385-417.
- Dauphin F, Ting V, Payette P, Dennis M & Hamel E. (1991). Vasocontractile muscarinic M1 receptors in cat cerebral arteries: pharmacological identification and detection of mRNA. *European journal of pharmacology: Molecular pharmacology* **207**, 319-327.
- de la Torre JC. (2002). Vascular basis of Alzheimer's pathogenesis. *Annals of the New York Academy of Sciences* **977**, 196-215.
- Delépine L & Aubineau P. (1997). Plasma protein extravasation induced in the rat dura mater by stimulation of the parasympathetic sphenopalatine ganglion. *Experimental Neurology* **147**, 389-400.

- Dickinson J & Paton JFR. (2012). *Cerebrovascular Hypertension*. Book Guild Ltd, Sussex, England.
- Duckles SP. (1981). Evidence for a functional cholinergic innervation of cerebral arteries. *Journal of pharmacology and experimental therapeutics* **217**, 544-548.
- Duckles SP & Said SI. (1982). Vasoactive intestinal peptide as a neurotransmitter in the cerebral circulation. *European journal of pharmacology* **78**, 371-374.
- Edvinsson L. (1987). Innervation of the cerebral circulation. *Annals of the New York Academy of Sciences* **519**, 334-348.
- Edvinsson L, Ekman R, Jansen I, Ottosson A & Uddman R. (1987). Peptide-containing nerve fibers in human cerebral arteries: Immunocytochemistry, radioimmunoassay and in vitro pharmacology. *Annals of Neurology* **21**, 431-437.
- Edvinsson L, Elsås T, Suzuki N, Shimizu T & Lee TJ. (2001). Origin and co-localization of nitric oxide synthase, CGRP, PACAP, and VIP in the cerebral circulation of the rat. *Microscopy research and technique* **53**, 221-228.
- Edvinsson L, Fahrenkrug J, Hanks J, Owman C, Sundler F & Uddman R. (1980). VIP (vasoactive intestinal polypeptide)-containing nerves of intracranial arteries in mammals. *Cell and tissue research* **208**, 135-142.
- Edvinsson L, Falck B & Owman C. (1977). Possibilities for a cholinergic action on smooth musculature and on sympathetic axons in brain vessels mediated by muscarinic and nicotinic receptors. *Journal of pharmacology and experimental therapeutics* **200**, 117-126.
- Edvinsson L, Gulbenkian S, Jansen I & Uddman R. (1994a). Neuropeptides in human cerebral arteries. In *The Human Brain Circulation*, ed. Bevan RD & Bevan JA, pp. 47-59. Humana Press, New York.
- Edvinsson L, Hara H & Uddman R. (1989). Retrograde tracing of nerve fibers to the rat middle cerebral artery with true blue: colocalization with different peptides. *Journal of cerebral blood flow and metabolism* **9**, 212-218.
- Edvinsson L, Jansen I, Sa MCE & Gulbenkian S. (1994b). Demonstration of neuropeptide containing nerves and vasomotor responses to perivascular peptides in human cerebral arteries. *Cephalalgia : an international journal of headache* **14**, 88-96.
- Edvinsson L, Nielsen KC, Owman C & Spörng B. (1972a). Cholinergic mechanisms in pial vessels. *Zellforsch* **134**, 311-325.

- Edvinsson L, Nielsen KC, Owman C & West KA. (1972b). Sympathetic neural influence on norepinephrine vasoconstriction in brain vessels. *Archives of neurology* **27**, 492-495.
- Edvinsson L, Owman C & Siesjö B. (1976a). Physiological role of cerebrovascular sympathetic nerves in the autoregulation of cerebral blood flow. *Brain research* **117**, 519-523.
- Edvinsson L, Owman C & Sjöberg N-O. (1976b). Autonomic nerves, mast cells, and amine receptors in human brain vessels. A histochemical and pharmacological study. *Brain research* **115**, 377-393.
- Elhusseiny A & Hamel E. (2000). Muscarinic-but not nicotinic-acetylcholine receptors mediate a nitric oxide-dependent dilation in brain cortical arterioles: A possible role for the M5 receptor subtype. *Journal of cerebral blood flow and metabolism* **20**, 298-305.
- Fahrenkrug J, Hannibal J, Tams J & Georg B. (2000). Immunohistochemical localization of the VIP1 receptor (VPAC1R) in rat cerebral blood vessels: relation to PACAP and VIP containing nerves. *Journal of cerebral blood flow and metabolism*.
- Fan N-C, Tsai C-M, Hsu C-N, Huang L-T & Tain Y-L. (2013). N-acetylcysteine prevents hypertension via regulation of the ADMA-DDAH pathway in young spontaneously hypertensive rats. *BioMed Research International* **2013**, 9.
- Fan Q & Smith PG. (1993). Decreased vasoactive intestinal polypeptide-immunoreactivity of parasympathetic neurons and target innervation following long-term sympathectomy. *Regulatory peptides* **48**, 337-343.
- Faraci FM & Brian JE. (1994). Nitric oxide and the cerebral circulation. *Stroke; a journal of cerebral circulation* **25**, 692-703.
- Faraci FM, Brian JE & Heistad DD. (1995). Response of cerebral blood vessels to an endogenous inhibitor of nitric oxide synthase. *American journal of physiology - Heart and circulatory physiology* **269**, H1522-H1527.
- Faraci FM & Heistad DD. (1990). Regulation of large cerebral arteries and cerebral microvascular pressure. *Circulation research* **66**, 8-17.
- Florence VM & Bevan JA. (1979). Biochemical determinations of cholinergic innervation in cerebral arteries. *Circulation research* **45**, 212-218.
- Frąckowiak H & Jakubowski H. (2008). Arterial vascularization in the giraffe brain. *Annales Zoologici Fennici* **45**, 353-359.

- Furchgott RF & Zawadzki JV. (1980). The obligatory role of endothelial cells in the relaxation of arterial smooth muscle by acetylcholine. *Nature* **288**, 373-376.
- Garcia-Villalon AL, Krause DN, Ehlert FJ & Duckles SP. (1991). Heterogeneity of muscarinic receptor subtypes in cerebral blood vessels. *Journal of pharmacology and experimental therapeutics* **258**, 304-310.
- Garthwaite J & Boulton C. (1995). Nitric oxide signaling in the central nervous system. *Annual review of physiology* **57**, 683-706.
- Gaw AJ, Aberdeen J, Humphrey PPA, Wadsworth RM & Burnstock G. (1991). Relaxation of sheep cerebral arteries by vasoactive intestinal polypeptide and neurogenic stimulation: inhibition by L-NG-monomethyl arginine in endothelium-denuded vessels. *British journal of pharmacology* **102**, 567-572.
- Ghia J-E, Blennerhassett P, El-Sharkawy RT & Collins SM. (2007). The protective effect of the vagus nerve in a murine model of chronic relapsing colitis. *American journal of physiology - Gastrointestinal and liver physiology* **293**, G711-G718.
- Gibbins IL, Brayden JE & Bevan JA. (1984a). Distribution and origins of VIP-immunoreactive nerves in the cephalic circulation of the cat. *Peptides* **5**, 209-212.
- Gibbins IL, Brayden JE & Bevan JA. (1984b). Perivascular nerves with immunoreactivity to vasoactive intestinal polypeptide in cephalic arteries of the cat: distribution, possible origins and functional implications. *Neuroscience* **13**, 1327-1346.
- Gibbins IL & Morris JL. (1988). Co-existence of immunoreactivity to neuropeptide Y and vasoactive intestinal peptide in non-noradrenergic axons innervating guinea pig cerebral arteries after sympathectomy. *Brain research* **444**, 402-406.
- Goadsby PJ. (2013). Chapter 16 - Autonomic nervous system control of the cerebral circulation. In *Handbook of Clinical Neurology*, ed. Ruud MB & Dick FS, pp. 193-201. Elsevier.
- Gonzalez C, Barroso C, Martin C, Gulbenkian S & Estrada C. (1997). Neuronal nitric oxide synthase activation by vasoactive intestinal peptide in bovine cerebral arteries. *Journal of cerebral blood flow and metabolism* **17**, 977-984.
- Grant S, Lutz EM, McPhaden AR & Wadsworth RM. (2005). Location and function of VPAC1, VPAC2 and NPR-C receptors in VIP-induced vasodilation of porcine basilar arteries. *Journal of cerebral blood flow and metabolism* **26**, 58-67.

- Greig JA, Shirley R, Graham D, Denby L, Dominiczak AF, Work LM & Baker AH. (2010). Vascular-Targeting Antioxidant Therapy in a Model of Hypertension and Stroke. *Journal of cardiovascular pharmacology* **56**, 642-650.
- Grimm MC, Newman R, Hassim Z, Cuan N, Connor SJ, Le Y, Wang JM, Oppenheim JJ & Lloyd AR. (2003). Cutting edge: vasoactive intestinal peptide acts as a potent suppressor of inflammation in vivo by trans-deactivating chemokine receptors. *Journal of immunology (Baltimore, Md : 1950)* **171**, 4990-4994.
- Gulbenkian S, Uddman R & Edvinsson L. (2001). Neuronal messengers in the human cerebral circulation. *Peptides* **22**, 995-1007.
- Hamel E. (2006). Perivascular nerves and the regulation of cerebrovascular tone. *Journal of applied physiology* **100**, 1059-1064.
- Hara H, Hamill GS & Jacobowitz DM. (1985). Origin of cholinergic nerves to the rat major cerebral arteries: Coexistence with vasoactive intestinal polypeptide. *Brain Research Bulletin* **14**, 179-188.
- Hara H, Jansen I, Ekman R, Hamel E, MacKenzie ET, Uddman R & Edvinsson L. (1989). Acetylcholine and vasoactive intestinal peptide in cerebral blood vessels: effect of extirpation of the sphenopalatine ganglion. *Journal of cerebral blood flow and metabolism* **9**, 204-211.
- Hara H & Kobayashi S. (1987). Vasoactive-intestinal-polypeptide (VIP)-like immunoreactive cells in the skull base of rats. *Histochemistry* **87**, 217-221.
- Hara H & Weir B. (1986). Pathway of acetylcholinesterase containing nerves to the major cerebral arteries in rats. *The Journal of comparative neurology* **250**, 245-252.
- Hara H, Zhang Q-J, Kuroyanagi T & Kobayashi S. (1993). Parasympathetic cerebrovascular innervation: an anterograde tracing from the sphenopalatine ganglion in the rat. *Neurosurgery* **32**, 822-827.
- Hardebo JE, Arbab M, Suzuki N & Svendgaard NA. (1991). Pathways of parasympathetic and sensory cerebrovascular nerves in monkeys. *Stroke; a journal of cerebral circulation* **22**, 331-342.
- Hardebo JE, Suzuki N, Ekblad E & Owman C. (1992). Vasoactive intestinal polypeptide and acetylcholine coexist with neuropeptide Y, dopamine-beta-hydroxylase, tyrosine hydroxylase, substance P or calcitonin gene-related peptide in neuronal subpopulations in cranial parasympathetic ganglia of rat. *Cell and tissue research* **267**, 291-300.
- Heistad DD & Marcus ML. (1978). Evidence that neural mechanisms do not have important effects on cerebral blood flow. *Circulation research* **42**, 295-302.

- Heistad DD, Marcus ML & Gross PM. (1978). Effects of sympathetic nerves on cerebral vessels in dog, cat, and monkey. *The American journal of physiology* **235**, H544-H552.
- Henninger N & Fisher M. (2007). Stimulating circle of Willis nerve fibers preserves the diffusion-perfusion mismatch in experimental stroke. *Stroke; a journal of cerebral circulation* **38**, 2779-2786.
- Hibbs Jr JB, Taintor RR, Vavrin Z & Rachlin EM. (1988). Nitric oxide: a cytotoxic activated macrophage effector molecule. *Biochemical and biophysical research communications* **157**, 87-94.
- Hoeger S, Bergstraesser C, Selhorst J, Fontana J, Birck R, Waldherr R, Beck G, Sticht C, Seelen MA, Van Son WJ, Leuvenink H, Ploeg R, Schnuelle P & Yard BA. (2010). Modulation of brain dead induced inflammation by vagus nerve stimulation. *American journal of transplantation* **10**, 477-489.
- Iadecola C, Beitz AJ, Renno W, Xu X, Mayer B & Zhang F. (1993). Nitric oxide synthase-containing neural processes on large cerebral arteries and cerebral microvessels. *Brain research* **606**, 148-155.
- Ignarro LJ. (1991). Signal transduction mechanisms involving nitric oxide. *Biochemical pharmacology* **41**, 485-490.
- Iwayama T, Furness J & Burnstock G. (1970). Dual adrenergic and cholinergic innervation of the cerebral arteries of the rat. An ultrastructural study. *Circulation research* **26**, 635-646.
- Kadota O, Matsuda S, Ohta S, Kumon Y, Sakaki S & Sakanaka M. (1996). Origins of nitric oxide synthase-containing nerve fibers in the rat basilar artery with reference to the fine structure of the nerve fibers. *Brain research* **706**, 129-136.
- Kano M, Moskowitz MA & Yokota M. (1991). Parasympathetic denervation of rat pial vessels significantly increases infarction volume following middle cerebral artery occlusion. *Journal of cerebral blood flow and metabolism* **11**, 628-637.
- Kashimoto K, Nagano Y, Suitani Y, Hamanaka K, Mizumoto T, Tomizaki K, Takahata H, Nagamoto A, Ohata A, Yoshihara S & Ichimura T. (1996). Structure-activity relationship studies of PACAP-27 and VIP analogues. *Annals of the New York Academy of Sciences* **805**, 505-510.
- Kawamura K & Takebayashi S. (1994). The development of noradrenaline-, acetylcholinesterase-, neuropeptide Y- and vasoactive intestinal polypeptide-containing nerves in human cerebral arteries. *Neuroscience letters* **175**, 1-4.

- Kawashima K, Watanabe N, Oohata H, Fujimoto K, Suzuki T, Ishizaki Y, Morita I & Murota S-i. (1990). Synthesis and release of acetylcholine by cultured bovine arterial endothelial cells. *Neuroscience letters* **119**, 156-158.
- Keller JT, Beduk A & Saunders MC. (1985). Origin of fibers innervating the basilar artery of the cat. *Neuroscience letters* **58**, 263-268.
- Khurana D, Kaul S, Bornstein NM & for the Imp ACTsg. (2009). Implant for augmentation of cerebral blood flow trial 1: a pilot study evaluating the safety and effectiveness of the Ischaemic Stroke System for treatment of acute ischaemic stroke. *International journal of stroke* **4**, 480-485.
- Kielstein JT, Donnerstag F, Gasper S, Menne J, Kielstein A, Martens-Lobenhoffer J, Scalera F, Cooke JP, Fliser D & Bode-Böger SM. (2006). ADMA increases arterial stiffness and decreases cerebral blood flow in humans. *Stroke; a journal of cerebral circulation* **37**, 2024-2029.
- Kielstein JT & Zoccali C. (2005). Asymmetric dimethylarginine: a cardiovascular risk factor and a uremic toxin coming of age? *American journal of kidney diseases* **46**, 186-202.
- Kimura T, Yu J-G, Edvinsson L & Lee TJF. (1997). Cholinergic, nitric oxidergic innervation in cerebral arteries of the cat. *Brain research* **773**, 117-124.
- Knutsson M & Edvinsson L. (2002). Distribution of mRNA for VIP and PACAP receptors in human cerebral arteries and cranial ganglia. *Neuroreport* **13**, 507-509.
- Kobayashi S, Kyoshima K, Olschowka JA & Jacobowitz DM. (1983). Vasoactive intestinal polypeptide immunoreactive and cholinergic nerves in the whole mount preparation of the major cerebral arteries of the rat. *Histochemistry* **79**, 377-381.
- Kobayashi S, Tsukahara S, Sugita K, Matsuo K & Nagata T. (1981). Histochemical studies on the postnatal development of autonomic nerves in mice cerebral arteries. *Histochemistry* **73**, 15-20.
- Koketsu N, Moskowitz MA, Kontos HA, Yokota M & Shimizu T. (1992). Chronic parasympathetic sectioning decreases regional cerebral blood flow during hemorrhagic hypotension and increases infarct size after middle cerebral artery occlusion in spontaneously hypertensive rats. *Journal of cerebral blood flow and metabolism* **12**, 613-620.
- Lawrence WE & Rewell RE. (1948). The cerebral blood supply in the giraffidae. *Proceedings of the Zoological Society of London* **118**, 202-212.

- Leceta J, Gomariz RP, Martinez C, Abad C, Ganea D & Delgado M. (2000). Receptors and Transcriptional Factors Involved in the Anti-Inflammatory Activity of VIP and PACAP. *Annals of the New York Academy of Sciences* **921**, 92-102.
- Lee TJ-F. (2000). Nitric oxide and the cerebral vascular function. *J Biomed Sci* **7**, 16-26.
- Lee TJ, Hume WR, Su C & Bevan JA. (1978). Neurogenic vasodilation of cat cerebral arteries. *Circulation research* **42**, 535-542.
- Lee TJ & Saito A. (1986). Altered cerebral vessel innervation in spontaneously hypertensive and renal hypertensive rats. *Journal of hypertension, Supplement* **4**, S201-203.
- Lee TJ, Su C & Bevan JA. (1976). Neurogenic sympathetic vasoconstriction of the rabbit basilar artery. *Circulation research* **39**, 120-126.
- Lee TJF. (1980). Direct evidence against acetylcholine as the dilator transmitter in the cat cerebral-artery. *European journal of pharmacology* **68**, 393-394.
- Lee TJF. (1982). Morphopharmacological study of cerebral vasodilator and constrictor nerves. *Developments in neuroscience* **14**, 431-439.
- Lee TJF. (2002). Adrenergic and cholinergic modulation of cerebrovascular nitrenergic vasodilation. *Brain activation and CBF control, Proceedings* **1235**, 337-345.
- Lee TJF, Chang HH, Lee HC, Chen PY, Lee YC, Kuo JS & Chen MF. (2011). Axo-axonal interaction in autonomic regulation of the cerebral circulation. *Acta physiologica* **203**, 25-35.
- Lee TJF, Fang YX & Nickols GA. (1990). Multiple transmitters are involved in cerebral neurogenic vasodilation. *Blood vessels* **27**, 43-43.
- Lee TJF, Liu J & Evans MS. (2001). Cholinergic-nitrenergic transmitter mechanisms in the cerebral circulation. *Microscopy research and technique* **53**, 119-128.
- Leiper J & Nandi M. (2011). The therapeutic potential of targeting endogenous inhibitors of nitric oxide synthesis. *Nature reviews drug discovery* **10**, 277-291.
- Levi H, Schoknecht K, Prager O, Chassidim Y, Weissberg I, Serlin Y & Friedman A. (2012). Stimulation of the sphenopalatine ganglion induces reperfusion and blood-brain barrier protection in the photothrombotic stroke model. *PLoS ONE* **7**, e39636.

- Linder S, Barkhem T, Norberg A, Persson H, Schalling M, Hökfelt T & Magnusson G. (1987). Structure and expression of the gene encoding the vasoactive intestinal peptide precursor. *Proceedings of the National Academy of Sciences* **84**, 605-609.
- Liu J, Evans MS, Brewer GJ & Lee TJF. (2000). N-type Ca^{2+} channels in cultured rat sphenopalatine ganglion neurons: an immunohistochemical and electrophysiological study. *Journal of cerebral blood flow and metabolism* **20**, 183-191.
- Liu J, Evans MS & Lee TJ. (2002). Presynaptic muscarinic $\text{M}(2)$ -receptor-mediated inhibition of N-type $\text{Ca}(2+)$ channels in cultured sphenopalatine ganglion: direct evidence for acetylcholine inhibition of cerebral nitroergic neurogenic vasodilation. *The Journal of pharmacology and experimental therapeutics* **302**, 397-405.
- Liu J & Lee TJ. (1999). Mechanism of prejunctional muscarinic receptor-mediated inhibition of neurogenic vasodilation in cerebral arteries. *American journal of physiology - Heart and circulatory physiology* **276**, H194-204.
- Luff SE & McLachlan EM. (1989). Frequency of neuromuscular junctions on arteries of different dimensions in the rabbit, guinea pig and rat. *Blood vessels* **26**, 95-106.
- Marina N, Ang R, Machhada A, Kasymov V, Karagiannis A, Hosford PS, Mosienko V, Teschemacher AG, Vihko P, Paton JFR, Kasparov S & Gourine AV. (2015). Brainstem hypoxia contributes to the development of hypertension in the spontaneously hypertensive rat. *Hypertension* **65**, 775-783.
- Martinez C, Abad C, Delgado M, Arranz A, Juarranz MG, Rodriguez-Henche N, Brabet P, Leceta J & Gomariz RP. (2002). Anti-inflammatory role in septic shock of pituitary adenylate cyclase-activating polypeptide receptor. *Proceedings of the National Academy of Sciences* **99**, 1053-1058.
- Matsuyama T, Shiosaka S, Wanaka A, Yoneda S, Kimura K, Hayakawa T, Emson PC & Tohyama M. (1985). Fine structure of peptidergic and catecholaminergic nerve fibers in the anterior cerebral artery and their interrelationship: An immunoelectron microscopic study. *The Journal of comparative neurology* **235**, 268-276.
- McCulloch J, Uddman R, Kingman TA & Edvinsson L. (1986). Calcitonin gene-related peptide: functional role in cerebrovascular regulation. *Proceedings of the National Academy of Sciences* **83**, 5731-5735.
- McDonald D & Potter J. (1951). The distribution of blood to the brain. *The journal of physiology* **114**, 356-371.

- McFarland WL, Jacobs MS & Morgane PJ. (1979). *Blood Supply to the Brain of the Dolphin, Tursiops Truncatus, with Comparative Observations on Special Aspects of the Cerebrovascular Supply of Other Vertebrates*, vol. 3. Ankho International.
- Miao FJ & Lee TJ. (1990). Cholinergic and VIPergic innervation in cerebral arteries: a sequential double-labeling immunohistochemical study. *Journal of cerebral blood flow and metabolism* **10**, 32-37.
- Miao FJP & Lee TJF. (1991). VIP-ergic and cholinergic innervations in internal carotid arteries of the cat and rat. *Journal of cardiovascular pharmacology* **18**, 369-378.
- Minami Y, Kimura H, Aimi Y & Vincent SR. (1994). Projections of nitric oxide synthase-containing fibers from the sphenopalatine ganglion to cerebral arteries in the rat. *Neuroscience* **60**, 745-759.
- Mione MC, Dhital KK, Amenta F & Burnstock G. (1988). An increase in the expression of neuropeptidergic vasodilator, but not vasoconstrictor, cerebrovascular nerves in aging rats. *Brain research* **460**, 103-113.
- Mione MC, Sancesario G, D'Angelo V & Bernardi G. (1991). Increase of dopamine β -hydroxylase immunoreactivity in non-noradrenergic nerves of rat cerebral arteries following long-term sympathectomy. *Neuroscience letters* **123**, 167-171.
- Mueller SM, Heistad DD & Marcus ML. (1977). Total and regional cerebral blood flow during hypotension, hypertension, and hypocapnia. Effect of sympathetic denervation in dogs. *Circulation research* **41**, 350-356.
- Murthy KS, Teng B-Q, Zhou H, Jin J-G, Grider JR & Makhlof GM. (2000). Gi-1/Gi-2-dependent signaling by single-transmembrane natriuretic peptide clearance receptor. *American journal of physiology - Gastrointestinal and liver physiology* **278**, G974-G980.
- Nagel EL, Morgane PJ, McFarland WL & Galllano RE. (1968). Rete mirabile of dolphin: its pressure-damping effect on cerebral circulation. *Science* **161**, 898-900.
- Nakai M, Tamaki K, Ogata J, Matsui Y & Maeda M. (1993). Parasympathetic cerebrovasodilator center of the facial nerve. *Circulation research* **72**, 470-475.
- Napoli C, Sica V, de Nigris F, Pignalosa O, Condorelli M, Ignarro LJ & Liguori A. (2004). Sulfhydryl angiotensin-converting enzyme inhibition induces sustained reduction of systemic oxidative stress and improves the nitric oxide pathway in patients with essential hypertension. *American heart journal* **148**, 172.

- Nozaki K, Moskowitz MA, Maynard KI, Koketsu N, Dawson TM, Bredt DS & Snyder SH. (1993). Possible origins and distribution of immunoreactive nitric oxide synthase-containing nerve fibers in cerebral arteries. *Journal of cerebral blood flow & metabolism* **13**, 70-79.
- Owman C. (2011). Neurogenic control of the vascular system: focus on cerebral circulation. In *Comprehensive physiology*, pp. 525-580. John Wiley & Sons, Inc.
- Owman C, Edvinsson L & Nielsen KC. (1974). Autonomic neuroreceptor mechanisms in brain vessels. *Journal of vascular research* **11**, 2-31.
- Parnavelas JG, Kelly W & Burnstock G. (1985). Ultrastructural localization of choline acetyltransferase in vascular endothelial cells in rat brain. *Nature* **316**, 724-725.
- Paton JF, Dickinson CJ & Mitchell G. (2009). Harvey Cushing and the regulation of blood pressure in giraffe, rat and man: introducing 'Cushing's mechanism'. *Experimental physiology* **94**, 11-17.
- Pavlov VA, Wang H, Czura CJ, Friedman SG & Tracey KJ. (2003). The cholinergic anti-inflammatory pathway: A missing link in neuroimmunomodulation. *Molecular medicine* **9**, 125-134.
- Phillips JK, Vidovic M & Hill CE. (1997). Variation in mRNA expression of alpha-adrenergic, neurokinin and muscarinic receptors amongst four arteries of the rat. *Journal of the autonomic nervous system* **62**, 85-93.
- Purves J. (1972). *The Physiology of the Cerebral Circulation*. University Press.
- Reale M, Iarlori C, Gambi F, Feliciani C, Salone A, Toma L, DeLuca G, Salvatore M, Conti P & Gambi D. (2004). Treatment with an acetylcholinesterase inhibitor in Alzheimer patients modulates the expression and production of the pro-inflammatory and anti-inflammatory cytokines. *Journal of neuroimmunology* **148**, 162-171.
- Rubino A & Burnstock G. (1996). Capsaicin-sensitive sensory-motor neurotransmission in the peripheral control of cardiovascular function. *Cardiovascular research* **31**, 467-479.
- Rush AJ, Sackeim HA, Marangell LB, George MS, Brannan SK, Davis SM, Lavori P, Howland R, Kling MA, Rittberg B, Carpenter L, Ninan P, Moreno F, Schwartz T, Conway C, Burke M & Barry JJ. (2005). Effects of 12 Months of Vagus Nerve Stimulation in Treatment-Resistant Depression: A Naturalistic Study. *Biological psychiatry* **58**, 355-363.
- Sadoshima S, Busija D, Brody M & Heistad D. (1981). Sympathetic nerves protect against stroke in stroke-prone hypertensive rats. A preliminary report. *Hypertension* **3**, 1124-127.

- Saito A, Wu JY & Lee TJ. (1985). Evidence for the presence of cholinergic nerves in cerebral arteries: an immunohistochemical demonstration of choline acetyltransferase. *Journal of cerebral blood flow and metabolism* **5**, 327-334.
- Santizo R, Baughman VL & Pelligrino DA. (2000). Relative contributions from neuronal and endothelial nitric oxide synthases to regional cerebral blood flow changes during forebrain ischemia in rats. *Neuroreport* **11**, 1549-1553.
- Schäfer MKH, Eiden LE & Weihe E. (1998). Cholinergic neurons and terminal fields revealed by immunohistochemistry for the vesicular acetylcholine transporter. II. The peripheral nervous system. *Neuroscience* **84**, 361-376.
- Schoen JC, Boysen MM, Warren CR, Chakravarthy B & Lotfipour S. (2011). Vertebrobasilar artery occlusion. *Western journal of emergency medicine* **12**, 233-239.
- Schuenke M, Schulte E, Schumacher U, Voll M & K W. (2010). *Head and Neuroanatomy (THIEME Atlas of anatomy)*. Thieme.
- Scremin OU. (2004). The Rat Nervous System. ed. Paxinos G, pp. 1165-1202. Elsevier Science.
- Seebeck J, Löwe M, Kruse M-L, Schmidt WE, Mehdorn HM, Ziegler A & Hempelmann RG. (2002). The vasorelaxant effect of pituitary adenylate cyclase activating polypeptide and vasoactive intestinal polypeptide in isolated rat basilar arteries is partially mediated by activation of nitrergic neurons. *Regulatory peptides* **107**, 115-123.
- Seki Y, Suzuki Y, Baskaya MK, Kano T, Saito K, Takayasu M, Shibuya M & Sugita K. (1995). The effects of pituitary adenylate cyclase-activating polypeptide on cerebral arteries and vertebral artery blood flow in anesthetized dogs. *European journal of pharmacology* **275**, 259-266.
- Seylaz J, Hara H, Pinard E, Mraovitch S, MacKenzie ET & Edvinsson L. (1988). Effect of stimulation of the sphenopalatine ganglion on cortical blood flow in the rat. *Journal of cerebral blood flow and metabolism* **8**, 875-878.
- Shalev A & Gross Y. (2010). Stimulation of the OTIC ganglion for treating medical conditions. Google Patents.
- Shimizu T. (1994). Distribution and pathway of the cerebrovascular nerve fibers from the otic ganglion in the rat: anterograde tracing study. *Journal of the autonomic nervous system* **49**, 47-54.
- Shimizu T, Rosenblum WI & Nelson GH. (1993). M3 and M1 receptors in cerebral arterioles in vivo: evidence for downregulated or ineffective M1 when endothelium is intact. *American journal of physiology - Heart and circulatory physiology* **264**, H665-H669.

- Si ML & Lee TJ. (2001). Presynaptic $\alpha 7$ -nicotinic acetylcholine receptors mediate nicotine-induced nitric oxidergic neurogenic vasodilation in porcine basilar arteries. *The Journal of pharmacology and experimental therapeutics* **298**, 122-128.
- Si ML & Lee TJF. (2002). $\alpha(7)$ -nicotinic acetylcholine receptors on cerebral perivascular sympathetic nerves mediate choline-induced nitrergic neurogenic vasodilation. *Circulation research* **91**, 62-69.
- Smith PG, Warn JD, Steinle JJ, Krizsan-Agbas D & Hasan W. (2002). Modulation of parasympathetic neuron phenotype and function by sympathetic innervation. *Autonomic neuroscience* **96**, 33-42.
- Solberg Y, Yarnitsky D, Borenstein N, Tanne D, Dayan A, Weiss S & Fisher M. (2008). Effectiveness of sphenopalatine ganglion stimulation therapy in focal ischemic stroke models: a 24-hour post-stroke window of treatment. In *Stroke; a journal of cerebral circulation*, pp. 665-665.
- Song J-N, Zhang M, Li D-D, Li M, An J-Y, Cheng M-F & Guo X-Y. (2014). Dynamic expression of the suppressor of cytokine signaling-3 and cytokines in the cerebral basilar artery of rats with subarachnoid hemorrhage, and the effect of acetylcholine. *Acta Neurochirurgica* **156**, 941-949.
- Spencer SE, Sawyer WB, Wada H, Platt KB & Loewy AD. (1990). CNS projections to the pterygopalatine parasympathetic preganglionic neurons in the rat: a retrograde transneuronal viral cell body labeling study. *Brain research* **534**, 149-169.
- Sundler F, Ekblad E, Hannibal J, Moller K, Zhang YZ, Mulder H, Elsås T, Grunditz T, Danielsen N, Fahrenkrug J & Uddman R. (1996). Pituitary adenylate cyclase-activating peptide in sensory and autonomic ganglia: localization and regulation. *Annals of the New York Academy of Sciences* **805**, 410-426.
- Suzuki N & Hardebo JE. (1991). The pathway of parasympathetic nerve fibers to cerebral vessels from the otic ganglion in the rat. *Journal of the autonomic nervous system* **36**, 39-46.
- Suzuki N & Hardebo JE. (1993). The cerebrovascular parasympathetic innervation. *Cerebrovasc Brain Metab Rev* **5**, 33-46.
- Suzuki N, Hardebo JE, Kahrstrom J & Owman C. (1990a). Neuropeptide Y co-exists with vasoactive intestinal polypeptide and acetylcholine in parasympathetic cerebrovascular nerves originating in the sphenopalatine, otic, and internal carotid ganglia of the rat. *Neuroscience* **36**, 507-519.
- Suzuki N, Hardebo JE, Kahrstrom J & Owman C. (1990b). Selective electrical stimulation of postganglionic cerebrovascular parasympathetic nerve fibers originating from the

- sphenopalatine ganglion enhances cortical blood flow in the rat. *Journal of cerebral blood flow and metabolism* **10**, 383-391.
- Suzuki N, Hardebo JE & Owman C. (1988). Origins and pathways of cerebrovascular vasoactive intestinal polypeptide-positive nerves in rat. *Journal of cerebral blood flow and metabolism* **8**, 697-712.
- Suzuki N, Hardebo JE & Owman C. (1990c). Origins and pathways of choline acetyltransferase-positive parasympathetic nerve fibers to cerebral vessels in rat. *Journal of cerebral blood flow and metabolism* **10**, 399-408.
- Suzuki Y, McMaster D, Lederis K & Rorstad OP. (1984). Characterization of the relaxant effects of vasoactive intestinal peptide (VIP) and PHI on isolated brain arteries. *Brain research* **322**, 9-16.
- Taktakishvili OM, Lin L-H, Vanderheyden AD, Nashelsky MB & Talman WT. (2010). Nitroxidergic innervation of human cerebral arteries. *Autonomic neuroscience* **156**, 152-153.
- Talman WT, Corr J, Nitschke Dragon D & Wang D. (2007). Parasympathetic stimulation elicits cerebral vasodilatation in rat. *Autonomic neuroscience* **133**, 153-157.
- Talman WT & Dragon DN. (2000). Parasympathetic nerves influence cerebral blood flow during hypertension in rat. *Brain research* **873**, 145-148.
- Tayebati SK, Di Tullio MA, Tomassoni D & Amenta F. (2003). Localization of the m5 muscarinic cholinergic receptor in rat circle of Willis and pial arteries. *Neuroscience* **122**, 205-211.
- Thompson BG, Pluta RM, Girton ME & Oldfield EH. (1996). Nitric oxide mediation of chemoregulation but not autoregulation of cerebral blood flow in primates. *Journal of neurosurgery* **84**, 71-78.
- Toda N. (1982). Relaxant responses to transmural stimulation and nicotine of dog and monkey cerebral arteries. *American journal of physiology* **243**, H145-H153.
- Toda N, Ayajiki K & Okamura T. (1997). Inhibition of nitroxidergic nerve function by neurogenic acetylcholine in monkey cerebral arteries. *The journal of physiology* **498**, 453-461.
- Toda N, Ayajiki K & Okamura T. (2009). Cerebral blood flow regulation by nitric oxide: recent advances. *Pharmacological reviews* **61**, 62-97.
- Toda N, Ayajiki K, Tanaka T & Okamura T. (2000a). Preganglionic and postganglionic neurons responsible for cerebral vasodilation mediated by nitric oxide in anesthetized dogs. *Journal of cerebral blood flow and metabolism* **20**, 700-708.

- Toda N, Ayajiki K, Yoshida K, Kimura H & Okamura T. (1993). Impairment by damage of the pterygopalatine ganglion of nitroxidergic vasodilator nerve function in canine cerebral and retinal arteries. *Circulation research* **72**, 206-213.
- Toda N & Okamura T. (2003). The pharmacology of nitric oxide in the peripheral nervous system of blood vessels. *Pharmacological reviews* **55**, 271-324.
- Toda N, Tanaka T, Ayajiki K & Okamura T. (2000b). Cerebral vasodilatation induced by stimulation of the pterygopalatine ganglion and greater petrosal nerve in anesthetized monkeys. *Neuroscience* **96**, 393-398.
- Toyoda K, Fujii K, Ibayashi S, Kitazono T, Nagao T, Takaba H & Fujishima M. (1998). Attenuation and recovery of brain stem autoregulation in spontaneously hypertensive rats. *Journal of cerebral blood flow and metabolism* **18**, 305-310.
- Tracey KJ. (2007). Physiology and immunology of the cholinergic antiinflammatory pathway. *The journal of clinical investigation* **117**, 289-296.
- Tsai C-M, Kuo H-C, Hsu C-N, Huang L-T & Tain Y-L. (2014). Metformin reduces asymmetric dimethylarginine and prevents hypertension in spontaneously hypertensive rats. *Translational research* **164**, 452-459.
- Tsai SH, Tew JM & Shipley MT. (1989). Cerebral arterial innervation: II. Development of calcitonin-gene-related peptide and norepinephrine in the rat. *The Journal of comparative neurology* **279**, 1-12.
- Tsai SH, Tew JM & Shipley MT. (1992). Development of cerebral arterial innervation: synchronous development of neuropeptide Y (NPY)- and vasoactive intestinal polypeptide (VIP)-containing fibers and some observations on growth cones. *Brain research Developmental brain research* **69**, 77-83.
- Tsukahara T, Usui H, Taniguchi T, Shimohama S, Fujiwara M & Handa H. (1986). Characterization of muscarinic cholinergic receptors in human and dog cerebral arteries. *Stroke; a journal of cerebral circulation* **17**, 300-305.
- Uddman R, Luts A, Håkanson R & Sundler F. (1993). Regulatory Peptides of the Parasympathetic Vascular Innervation. In *Vascular Innervation and Receptor Mechanisms: New Perspectives*, ed. Edvinsson L & Uddman R, pp. 203-220. Academic Press, San Diego, CA.
- Uddman R, Tajti J, Möller S, Sundler F & Edvinsson L. (1999). Neuronal messengers and peptide receptors in the human sphenopalatine and otic ganglia. *Brain research* **826**, 193-199.

- Vallance P, Leone A, Calver A, Collier J & Moncada S. (1992). Accumulation of an endogenous inhibitor of nitric oxide synthesis in chronic renal failure. *The Lancet* **339**, 572-575.
- van Charldorp K & van Zwieten P. (1989). Comparison of the muscarinic receptors in the coronary artery, cerebral artery and atrium of the pig. *Naunyn-Schmiedeberg's archives of pharmacology* **339**, 403-408.
- van Maanen MA, Lebre MC, van der Poll T, LaRosa GJ, Elbaum D, Vervoordeldonk MJ & Tak PP. (2009). Stimulation of nicotinic acetylcholine receptors attenuates collagen-induced arthritis in mice. *Arthritis and rheumatism* **60**, 114-122.
- van Westerloo DJ, Giebelen IA, Florquin S, Bruno MJ, Larosa GJ, Ulloa L, Tracey KJ & van der Poll T. (2006). The vagus nerve and nicotinic receptors modulate experimental pancreatitis severity in mice. *Gastroenterology* **130**, 1822-1830.
- Wahl M & Schilling L. (1993). Regulation of Cerebral Blood Flow — A Brief Review. In *Monitoring of Cerebral Blood Flow and Metabolism in Intensive Care*, ed. Unterberg A, Schneider G-H & Lanksch W, pp. 3-10. Springer, Vienna.
- Waki H, Gouraud SS, Maeda M & Paton JF. (2010). Evidence of specific inflammatory condition in nucleus tractus solitarii of spontaneously hypertensive rats. *Experimental physiology* **95**, 595-600.
- Waki H, Hendy EB, Hindmarch CCT, Gouraud S, Toward M, Kasparov S, Murphy D & Paton JFR. (2013). Excessive Leukotriene B4 in Nucleus Tractus Solitarii Is Prohypertensive in Spontaneously Hypertensive Rats. *Hypertension* **61**, 194-201.
- Waki H, Liu B, Miyake M, Katahira K, Murphy D, Kasparov S & Paton JF. (2007). Junctional adhesion molecule-1 is upregulated in spontaneously hypertensive rats: evidence for a prohypertensive role within the brain stem. *Hypertension* **49**, 1321-1327.
- Walters BB, Gillespie SA & Moskowitz MA. (1986). Cerebrovascular projections from the sphenopalatine and otic ganglia to the middle cerebral artery of the cat. *Stroke; a journal of cerebral circulation* **17**, 488-494.
- Wang H, Liao H, Ochani M, Justiniani M, Lin X, Yang L, Al-Abed Y, Wang H, Metz C, Miller EJ, Tracey KJ & Ulloa L. (2004). Cholinergic agonists inhibit HMGB1 release and improve survival in experimental sepsis. *Nature Medicine* **10**, 1216-1221.
- Wang H, Yu M, Ochani M, Amella CA, Tanovic M, Susarla S, Li JH, Wang H, Yang H, Ulloa L, Al-Abed Y, Czura CJ & Tracey KJ. (2003). Nicotinic acetylcholine receptor [alpha]7 subunit is an essential regulator of inflammation. *Nature* **421**, 384-388.

- Warren JB, Donnelly LE, Cullen S, Robertson BE, Ghatei MA, Bloom SR & MacDermot J. (1991). Pituitary adenylate cyclase-activating polypeptide: a novel, long-lasting, endothelium-independent vasorelaxant. *European journal of pharmacology* **197**, 131-134.
- White RP. (1987). Responses of human basilar arteries to vasoactive intestinal polypeptide. *Life sciences* **41**, 1155-1163.
- Willmot M, Gray L, Gibson C, Murphy S & Bath PMW. (2005). A systematic review of nitric oxide donors and l-arginine in experimental stroke; effects on infarct size and cerebral blood flow. *Nitric oxide* **12**, 141-149.
- Yamada H, Honda T, Yaginuma H, Kikuchi S & Sugiura Y. (2001a). Comparison of sensory and sympathetic innervation of the dura mater and posterior longitudinal ligament in the cervical spine after removal of the stellate ganglion. *The journal of comparative neurology* **434**, 86-100.
- Yamada M, Lamping KG, Duttaroy A, Zhang W, Cui Y, Bymaster FP, McKinzie DL, Felder CC, Deng C-X, Faraci FM & Wess J. (2001b). Cholinergic dilation of cerebral blood vessels is abolished in M5 muscarinic acetylcholine receptor knockout mice. *Proceedings of the National Academy of Sciences* **98**, 14096-14101.
- Yarnitsky D, Goor-Aryeh I, Bajwa ZH, Ransil BI, Cutrer FM, Sottile A & Burstein R. (2003). 2003 Wolff Award: Possible parasympathetic contributions to peripheral and central sensitization during migraine. *Headache: The journal of head and face pain* **43**, 704-714.
- Yarnitsky D, Gross Y, Lorian A, Shalev A, Lamensdorf I, Bornstein R, Shorer S, Mayevsky A, Patel KP, Abbott NJ & Mayhan WG. (2004a). Blood-brain barrier opened by stimulation of the parasympathetic sphenopalatine ganglion: a new method for macromolecule delivery to the brain. *Journal of neurosurgery* **101**, 303-309.
- Yarnitsky D, Gross Y, Lorian A, Shalev A, Shorer S, Tanaka T, Ayajiki K, Fujimiya M & Okamura T. (2004b). Increased BBB permeability by parasympathetic sphenopalatine ganglion stimulation in dogs. *Brain research* **1018**, 236-240.
- Yarnitsky D, Lorian A, Shalev A, Zhang Z-D, Takahashi M, Agbaje-Williams M & Macdonald RL. (2005). Reversal of cerebral vasospasm by sphenopalatine ganglion stimulation in a dog model of subarachnoid hemorrhage. *Surgical neurology* **64**, 5-11.
- Yoo J-H & Lee S-C. (2001). Elevated levels of plasma homocyst(e)ine and asymmetric dimethylarginine in elderly patients with stroke. *Atherosclerosis* **158**, 425-430.
- Yoshida K, Okamura T, Kimura H, Bredt DS, Snyder SH & Toda N. (1993). Nitric oxide synthase-immunoreactive nerve fibers in dog cerebral and peripheral arteries. *Brain research* **629**, 67-72.

- Yu J-G, Kimura T, Chang X-F & Lee TJF. (1998). Segregation of VIPergic–nitric oxidergic and cholinergic–nitric oxidergic innervation in porcine middle cerebral arteries. *Brain research* **801**, 78-87.
- Zhang E-T, Mikkelsen JD, Fahrenkrug J, Møller M, Kronborg D & Lauritzen M. (1991). Prepro-vasoactive intestinal polypeptide-derived peptide sequences in cerebral blood vessels of rats: on the functional anatomy of metabolic autoregulation. *Journal of cerebral bloodflow and metabolism* **11**, 932-938.
- Zhang W, Edvinsson L & Lee TJ. (1998a). Mechanism of nicotine-induced relaxation in the porcine basilar artery. *The Journal of pharmacology and experimental therapeutics* **284**, 790-797.
- Zhang W, Edvinsson L & Lee TJF. (1998b). Nicotine-induced NO-mediated cerebral neurogenic vasodilation is dependent on intact adrenergic innervation. *Naunyn-Schmiedeberg's archives of pharmacology* **358**, R91-R91.
- Zubcevic J, Waki H, Raizada MK & Paton JFR. (2011). Autonomic-immune-vascular interaction: An emerging concept for neurogenic hypertension. *Hypertension* **57**, 1026-1033.